

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 May 2001 (31.05.2001)

PCT

(10) International Publication Number
WO 01/38579 A2

(51) International Patent Classification⁷: **C12Q 1/68**,
C12N 5/10, 15/63, C07K 14/47, 16/18, C07H 21/04

(21) International Application Number: PCT/US00/32049

(22) International Filing Date:
21 November 2000 (21.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/166,923 22 November 1999 (22.11.1999) US
60/183,531 18 February 2000 (18.02.2000) US
09/717,321 20 November 2000 (20.11.2000) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US	60/166,923 (CIP)
Filed on	22 November 1999 (22.11.1999)
US	60/183,531 (CIP)
Filed on	18 February 2000 (18.02.2000)
US	09/717,321 (CIP)
Filed on	20 November 2000 (20.11.2000)

(71) Applicant (for all designated States except US): **CURAGEN CORPORATION** [US/US]; 555 Long Wharf Drive, 11th floor, New Haven, CT 06511 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GOULD-ROTHBERG, Bonnie, E.** [CA/US]; 1701 Moose Hill Road,

Guilford, CT 06437 (US). **DIPIPPO, Vincent, A.** [US/US]; 156 North Atwater Street, East Haven, CT 06512 (US). **RAMSEH, Tennore, M.** [IN/US]; 7 Sandpiper Lane, New Milford, CT 06776 (US). **GERWEIN, Robert, W.** [US/US]; 6 Montoya Circle, Brandford, CT 06405 (US).

(74) Agent: **ELRIFI, Ivor, R.**; Mintz, Levin, Cohn, Ferris, Glovsky and Popeo, P.C., One Financial Center, Boston, MA 02111 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD OF IDENTIFYING TOXIC AGENTS USING NSAID-INDUCED DIFFERENTIAL GENE EXPRESSION IN LIVER

(57) Abstract: The invention provides methods of identifying toxic agents, e.g., hepatotoxic agents, using differential gene expression. Also provided are methods of predicting the risk level and or injury type of NSAIDs. Also disclosed are novel nucleic acid sequences whose expression is differentially regulated by NSAIDs.

WO 01/38579 A2



METHOD OF IDENTIFYING TOXIC AGENTS USING NSAID-INDUCED DIFFERENTIAL GENE EXPRESSION IN LIVER

FIELD OF THE INVENTION

The invention relates generally to nucleic acids and polypeptides, and more particularly
5 to the identification of differentially expressed nucleic acids and proteins in liver.

BACKGROUND OF THE INVENTION

Liver is the primary organ for biotransformation of chemical compounds and their
detoxification. Liver injury produced by chemicals has been recognized for over 100 years,
and hepatic damage is one of the most common toxicities among drugs at pre-clinical and
10 clinical stages of drug development. Over 30% of new chemical entities (NCE) are generally
terminated due to adverse liver effects in humans. During a period of 30 years, hepatotoxicity
has been the major cause of drug withdrawal for safety reasons at the marketing stage,
accounting for 18% overall drug withdrawal. Many of the drugs that are withdrawn from
market due to hepatotoxicity produce lethality in a small percentage of patient population and
15 are classified as type II lesion (or idiosyncratic, sporadic) toxicity.

Non-steroidal anti-inflammatory drugs (NSAIDs) are a group of unrelated chemical
compounds that have been used to successfully treat rheumatic and musculoskeletal disease.
Unfortunately, unwanted hepatotoxic side effects have led to the premature market
withdrawal of several NSAIDs, including Cincophen, Benoxaprofen, Piroxicam, Suprofen,
20 and Bromfenac. The pervasiveness of idiosyncratic reactions of many NSAIDs has led the
Food and Drug Administration's Arthritis Advisory Board to conclude that NSAIDs as a group
should be considered to induce hepatotoxicity.

It is estimated that annual NSAIDs consumption in the U.S. exceeds 10,000 tons. Due
to this large consumption of NSAIDs for a wide variety of pain and inflammatory conditions,
25 it has become an important class of drugs responsible for liver injury, despite the overall
extremely low incidence of producing hepato-toxicity. Liver injury resulting from NSAIDs
can have several forms, including acute toxicity resulting from hepatocellular (parenchymal)
damage (e.g. necrosis) and arrested bile flow (cholestasis). The general mechanism that is
thought to mediate NSAIDs toxicity is an idiosyncratic reaction (Type II) to the drug (both
30 immunologic and metabolic), which is dose independent, and presumably results from
interindividual variation in drug metabolism. Currently no clear mechanism of drug-induced

idiosyncratic toxicity is available. Accordingly, there remains a great need to elucidate the molecular basis of idiosyncratic hepatotoxicity, such as NSAID-induced toxicity, including the identification of genes and proteins differentially expressed in response to administration of such drugs.

5

SUMMARY OF THE INVENTION

In accordance with the present invention, there are provided methods of screening and identifying test agents which induce hepatotoxicity, *e.g.* idiosyncratic hepatotoxicity. The methods of the invention are based in part on the discovery that certain nucleic acids are differentially expressed in liver tissue of animals treated with NSAIDs. These differentially
10 expressed nucleic acids include novel sequences that, while previously described, have not heretofore been identified as responsive to drugs, such as NSAIDs, which induce idiosyncratic hepatotoxicity.

In various aspects, the invention includes a method of screening a test agent for toxicity, *e.g.*, idiosyncratic hepatotoxicity. For example, in one aspect, the invention provides
15 a method of identifying a hepatotoxic agent by providing a test cell population comprising a cell capable of expressing one or more nucleic acids sequences responsive to drugs, *e.g.* NSAIDs, which induce idiosyncratic hepatotoxicity, contacting the test cell population with the test agent and comparing the expression of the nucleic acids sequences in the test cell population to the expression of the nucleic acids sequences in a reference cell population. An
20 alteration in expression of the nucleic acids sequences in the test cell population compared to the expression of the gene in the reference cell population indicates that the test agent is hepatotoxic. In one aspect, expression in the test cell population is compared to the expression of a reference cell population exposed to a NSAID that is classified as low risk, very low risk, or overdose risk of hepatotoxicity, thereby to predict whether the test agent has low, very low, or
25 overdose risk of hepatotoxicity. In another aspect, the test cell population is compared to the expression of a reference cell population exposed to a NSAID which induces a known type of hepatic injury, *e.g.* hepatocellular damage, cholestasis, or elevated transaminase level, thereby to predict whether the test agent is likely to induce a given type of hepatotoxic injury.

In a further aspect, the invention provides a method of assessing the hepatotoxicity,
30 *e.g.* idiosyncratic hepatotoxicity, of a test agent in a subject. The method includes providing from the subject a cell population comprising a cell capable of expressing one or more NSAID-responsive genes, and comparing the expression of the nucleic acids sequences to the

expression of the nucleic acids sequences in a reference cell population that includes cells from a subject whose exposure status to a hepatotoxic agent is known. An alteration in expression of the in the test cell population compared to the expression of the nucleic acids sequences in the reference cell population indicates the hepatotoxicity of the test agent in the subject.

Also provided are novel nucleic acids, as well as their encoded polypeptides, whose expression is responsive to the effects of NSAIDS.

Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

DETAILED DESCRIPTION OF THE INVENTION

The invention is based in part on the discovery of nucleic acid sequences which are differentially expressed in rodent liver cells upon administration of NSAIDS. The discovery includes groups of nucleic acid sequences whose expression is correlated with hepatotoxicity risk associated with, and injury type induced by, NSAID administration.

The differentially expressed nucleic acid sequences were identified by examining 29 different NSAIDS that have varying degrees of hepatotoxicity. These 29 drugs, shown in Table 1, below, were first categorized as low dose (1-75 mg/kg) and high dose (above 75 mg/kg) drugs, then classified as non-toxic, toxic, and those withdrawn from market (within each dose). Each of the 29 NSAIDS was administered orally to groups (3 animals per group) of 12 week old male Sprague Dawley rats for 72 hours (3 days) at the dosages specified in Table 1 (e.g. Naproxen: 54 mg/kg/day PO in QD x 3 days in H₂O). Vehicle controls (water, ethanol, canola oil) were also included (3 animals per group). The animals were sacrificed 24 hours after the final dose, liver tissue was removed on necroscopy, and total RNA was recovered from the dissected tissue. Complementary DNA (cDNA) was prepared and samples were processed through GENEALLING™ differential expression analysis, as described in U.S. Patent No. 5,871,697 and in Shimkets *et al.*, *Nature Biotechnology* 17: 798-803 (1999), the disclosures of which are hereby incorporated by reference herein.

Table 1: NSAIDS and Dosages Administered

Compound	Dose	Vehicle
<i>Acetaminophen</i>	133 mg/kg/day p.o. in QD x 3 days.	10 %Ethanol Vehicle
<i>Acetylsalicylic Acid</i>	200 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Benoxaprofen</i>	16 mg/kg/day p.o. in QD x 3 days.	H2O Vehicle
<i>Bromfenac</i>	7.5 mg/kg/day p.o. in QD x 3 days.	H2O Vehicle
<i>Celecoxib</i>	89 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Diclofenac</i>	38 mg/kg/day p.o. in QD x 3 days.	H2O Vehicle
<i>Etodolac</i>	30 mg/kg/day p.o. in QD x 3 days.	10% Ethanol Vehicle
<i>Felbinac</i>	33 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Fenoprofen</i>	154 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Flurbiprofen</i>	10 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Ibuprofen</i>	211 mg/kg/day p.o. in QD x 3 days.	H2O Vehicle
<i>Indomethacin</i>	4 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Ketoprofen</i>	10 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Ketorolac</i>	1.5 mg/kg/day p.o. in QD x 3 days.	10% Ethanol Vehicle
<i>Meclofenamate</i>	20 mg/kg/day p.o.in QD x 3 days.	H2O Vehicle
<i>Mefenamic Acid</i>	79 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Nabumetone</i>	143 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Naproxen</i>	54 mg/kg/day p.o. in QD x 3 days.	10% Ethanol Vehicle
<i>Olsalazine</i>	222 mg/kg/day p.o. in QD x 3 days.	H2O Vehicle
<i>Oxaprozin</i>	100 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Phenacetin</i>	100 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Phenylbutazone</i>	100 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Piroxicam</i>	20 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Sulindac</i>	77 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Sulphasalazine</i>	338 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Suprofen</i>	20 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Tenoxicam</i>	10 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle
<i>Tolmentin</i>	100 mg/kg/day p.o. in QD x 3 days.	H2O Vehicle
<i>Zomepirac</i>	19 mg/kg/day p.o. in QD x 3 days.	Canola Oil Vehicle

3635 gene fragments were initially found to be differentially expressed in rat liver tissue (analysis of variance, $p < 0.01$) in response to these compounds. The compounds were then classified according to hepatotoxicity risk, as indicated in Table 2.

Table 2: Hepatotoxicity Risk of NSAIDs

Compound	Risk
Acetaminophen	Overdose Risk
Acetylsalicylic Acid	Overdose Risk
Benoxaprofen	Low Risk
Bromfenac	Low Risk
Celecoxib	Unknown
Diclofenac	Low Risk
Etodolac	Very Low Risk
Felbinac	Unknown
Fenoprofen	Very Low Risk
Flurbiprofen	Very Low Risk
Ibuprofen	Very Low Risk
Indomethacin	Very Low Risk
Ketoprofen	Very Low Risk
Ketorolac	Unknown
Meclofenamate	Very Low Risk
Mefenamic Acid	Very Low Risk
Nabumetone	Very Low Risk
Naproxen	Very Low Risk
Olsalazine	Unknown
Oxaprozin	Very Low Risk
Phenacetin	Overdose Risk
Phenylbutazone	Low Risk
Piroxicam	Very Low Risk
Sulindac	Low Risk
Sulphasalazine	Unknown
Suprofen	Very Low Risk
Tenoxicam	Very Low Risk
Tolmentin	Very Low Risk
Zomepirac	Very Low Risk

In order to discriminate among these groups, the above compound set was divided into a training set (consisting of three compounds per group), and a test set (consisting of the remainder. This was done to minimize the reliance on the assumptions required for parametric analyses. Compounds with unknown risk were not used in this analysis. The training set employed is shown in Table 3.

Table 3: Training Set of NSAIDs by Risk Classification

Control	Low Risk	Very Low Risk	Overdose Risk
Sterile water	Benoxaprofen	Flurbiprofen	Acetaminophen
10% Ethanol	Phenylbutazone	Oxaprozin	Acetylsalicylic Acid
Canola oil	Sulindac	Tenoxicam	Phenacetin

The 3635 differentially expressed nucleic acid fragments were then analyzed using a stepwise multivariate analysis of variance as follows:

1. Calculate 3635 T2 (yi1) (Hoettelling's trace, one of the test statistics used for this analysis) values, one for each differentially expressed fragment. The fragment with the largest individual T2 value is selected as the first discriminatory set (yi1).
2. Calculate 3634 T2 (yi1,yi2) values, one for each combination of two fragments. The fragment pair with the largest individual T2 value are selected as the second discriminatory set.
3. Calculate 3626 T2 (yi1,yi2,yi3,...yi,10), one for each combination of ten fragments. The fragment set with the largest T2 value are selected as the final discriminatory set.

This stepwise procedure is used whenever the number of dependent variables (gene fragments) exceeds the number of independent variables (samples). In addition to fragment addition, fragment elimination occurs whenever an added fragment no longer contributes significant discriminatory power to the existing set. This eliminates bias as to the order fragments enter the model (Ahrens and Lauter, Mehrdimensionale Varianzanalyse, Akademie-Verlag, Berlin (1974); Dziuda, *Medical Inform.* 15(4): 319 (1990)).

This analysis protocol identified ten fragments that significantly ($p=6.02 \times 10^{-28}$) discriminated among the drugs in the test set. Two fragments on this list were not required to maintain the discriminatory ability and were subsequently removed ($p=3.96 \times 10^{-26}$). Differential expression of these gene fragments were successfully confirmed using an unlabeled oligonucleotide competition assay (Shimkets *et al.*, *Nature Biotechnology* 17: 198-803 (1999)). The 8 fragments (RISKMARKER 1-8) represent both novel and known rat genes for which the sequence identity to genes in public databases is either high (>90%), moderate (70-90%), or low (<70%).

The identity of these 8 hepatotoxicity risk discriminatory nucleic acid sequences (with GenBank accession numbers) are further described below. Where appropriate, the cloned sequence from isolation is provided; this sequence was then extended using either Genbank rat ESTs or from internally (Curagen Corporation) sequenced rat fragments. The extended contig sequence is provided as "consensus." Finally, the best BlastN and BlastX results are also provided. In some instances the cloned sequence is identical to a known rat gene, in those

RISKMARKER 1

10 The cloned sequence was assembled into a contig resulting in the following consensus
sequence:

1 TTTTTTTTTTTTTTTTCAAGTTCCAAAGACATTTTTTTTTTTTTTTATGATTCAAGGATTTATTAAGTCATACATGC
81 AAAACATACTGCTAACTGCATTAGCAAAAAGATCAATGTAAAAACTCCACAATTCTGCAACTGTCAATTGAAAAAAGTT
161 TGTTCTAGTGGTCGAAAGGCCAACACTGTGTTCTTGCCAGTGAGTTAGGTTGTACAGAACGGCGTTAGCACTAGCGCTT
241 GACAGAACC TCAGACCCAAAGGTACCGGAAGCATGTGTCCGCTGGGTGAGGTCTAGAGGGGGCGGCATCAATCACAT
321 GACAGTGTGGTACTCTGGCAAGACAGTGATGTTTCAGAATATCTAAAATAGTTTAAAAACTGTAAAGCCGCAGCAGCTG
401 ATTTCTACACCCAGTTACTAGAAAACGAAGGGAAGCACTAGTCAGCTGAGTAAAGGAAGGTGAAAACAGGAACGCACTTC
481 TACTATCTACCAAAAAAATCTCGAATGCATTATCAGAAAGATCTTATAGTACAGGT CAGACATATTGCTCGTTAAGAAAG
561 GGGGTCTTAAGAAAAAGCACTTGCTAAGTTAGCAACTGTGAGGATGGCCAGTTTAAATATGGACTCAACGCCCATCTGG
641 GGAGGGACAGCAGGGGGGAAGGGGGGCTCAAGAGAGACACTGATAAGATCGGCCATTTGT CATCTACTGTTTGACAGAAAT
721 TAACCGTTAAAAAGCTTTACCCGTGACACTTTTATTCA GTTGAATTACTCCATGTACAATGTAGTGTAAATTAATCTCTA
801 CTT CATATTAGTCAAAATACTGTCTGTCTCCTTTGATGACGTCGTGTTTCACACACTCCACCCAGCACCCCAGACTAG
881 GAACAGAATACTTCGTTAGAGGCAACACAGGAGCCAGAGTTCTGTTCAAAGCCTGCAGAAGCCGGTCAGCTGGTATTTTA
961 GAGAACTCACTATGAAATCAAAGAGCAGAGCTGTACACCCATCGTGACGTACAGTACAAAGTTACGTAATGAGCATGGG
1041 CTGATAAGTTACAGGTGCGTTACATGGCAGCGTGTCATT AAGGAGCGTGTGCTGTGTGCACACGGTCTGGGAGCTACGGGA
1121 GGGTCTGCACCCTGAGCCAGAAGCTGCAGTCTTCTTAAGGACAAAGTCTCTCAACAGCTTAGTGTTACGTGTTCTCA
GCACAACGCAACTAGTTCAACAAGGTATTTTGGCAATCTTAACTGAGCAAGAATAGGGGATTTT
1201

(SEQ ID NO: 2)

>gb:GENBANK-ID:HSA132695|acc:AJ132695 Homo sapiens rac1 gene - Homo sapiens, 28567 bp.

Minus Strand HSPs:

Query: 1146 GCTTCTGGGCTCAGG-GGTGCAGACCCCTCCCGTAGCTCCCAGACCGTGTGACACAGCACA 1088
||||||| | ||| ||||||||| || | | |||| | || | | |||| |

Query: 1087 GCCTCCTTAATGACACGCTGCC--ATGTAACGCA-CCTGT-AACTTATCAGCCCAT--GC 1034
||| | | | | | | | | | | | | | | | | | | | | |
5 Sbjct: 27041 GCGTAAAGCAGAACAGCTCCCGAATGAAGCGTTGCCATTGAACCTACCAGTGAGTTAGC 27100

Query: 1033 T-CA--T-TACGTAACCTTTGTACTGTAC-GTCACGATGG-GTGTAACAGCTCTGCTCTTT' 980
|| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27101 AGCACGTGTTCCCGACATAACATTGTACTGT-A--ATGGAGTG-AGCGTAGCAGCTCAGC 27156

10 Query: 979 GATTTCATAGTGAGT-TCTCTAAATACCAGCTGACCGGCTTCTG-CAGGCTTT-GAACA 923
||| | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27157 TCTTTGG-A-TCAGTCTTTGTGATTTTCATAGC-GA--GTTTCTGACCAGCTTTTGC GGA 27211

15 Query: 922 GAACTCTGG-CTC--CTG-TGTTGCCTCTAACGAAGTATTCTGTTCTAGTCGTGGGTGT 867
|| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27212 GATT-TGAACAGAAGTCTATTTCCTCTAATGAAGAATTCTGTT--TAGCTGTGGGTGT 27268

20 Query: 866 GCTGGGTGGAGTGTGTGAACACGACGCTCATCAAGGAGACAGACAGTATTTTGACTAA- 808
|| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27269 GCCGGGTGGGTGTGTGT-----GA-----TCAAGGACAAAGACAGTATTTTGACAAAA 27318

25 Query: 807 TATGAAGTAGAGATTAATT-TACACT--ACATTGTACATGGAGTAAT-TCAACTGAATAA 752
|| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27319 TACGAAGTGGAGATTACACTACATTGTACAAGG-A-ATGAA--AGTGTACACGGGTA-AA 27373

30 Query: 751 AAGTGTACACGGGTAAAGCTTTTT-AACGGTTAATTTCTG-TCAAACAGTAGATGACAA-A 695
|| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27374 AACTCTAAAAGTTAATTTCTGTCAAATGC-AGTAGATGATGAAGAAAGGTTGGTATTA 27432

35 Query: 694 TG-GCCGATCTTATCAGTGTCTCTCTTGAGCCCCCTTCCCCCTGCTGTCCTCCCCAGA 636
| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27433 TCAGGAAATGTTTCTTAAAGCT-T-TTCCTTCTC-TTACACCTGCCATGCCCTCCCCAAA 27489

40 Query: 635 TGGGGCGTTGAGTCCATATTTAAACTGGCCATCCTCACAGTTGCTAACTTAGCAAGTGCT 576
| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27490 TTGGGCATTAAATTCATCTTTAAACTGGTTGTTCTGTTAGTCGCTAACTTAGTAAGTGCT 27549

45 Query: 575 TTTCTT-TAGGACCCCTTCTTAACGAGCAATATGTCTGACCTGTACTATAAGATCTTTC 517
| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27550 TTTCTTATAGAACCCC-TTCTGACTGAGCAATATGCCTC-CTTGATTATAAATCTTTC 27607

50 Query: 516 TGATAATGCATTTCGAGATTTTTTTGGTAGATAGTAGAAGTGCCTTCC-TGTTTTACCT 458
| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27608 TGATAATGCATTAGAAGGTTTTTTGTGCGATTAGTAAAAGTGCTTTCCATGTTACTTTAT 27667

55 Query: 457 TCCTT--TACTCAG--CTGAC-T-AGTGCTTCCCTTCGTTT--TCTAGTAA-C-TGGGT- 409
T| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27668 TCAGAGCTAATAAGTGCTTTCCTTAGTT-TTCTAGTAAGTAGGTGAAAAATCATGTGTT 27726

60 Query: 408 GTAGAAATCACGTGCTGCGGCT-TTACAG-T--TT-TTAACTATTTTAGATATCTGAA 354
G | G T GT T T TT | G T TT | | | | | | | | | | | |
Sbjct: 27727 GCAGCTTTATAGTTTTTAAATATTTTAGATAATCTTAACTATG--A-ACCTTCTTAA 27783

65 Query: 353 ACATCACTGTCTTGCCAGAGTACCAACACTGTCATGTGATTGATGCCGCCCTCTAGAC 294
| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27784 -CATCACTGTCTTGCCAGATTACCGACACTGTCACCTGACCAATACTGACCC-TCCTTAC 27841

70 Query: 293 CTCACCCACGCGGACACATGCTTCCGGTA--C-CTTTGGGTCTGTGAGGTTT 245
| T | | | | | G | G | | | | | G | T | | GT | | | TTTG T T TG | GTT |
Sbjct: 27842 CTCGCCCACGCGGACACACGCTCCTGTAGTCGCTTTGCCTAT-TGATGTTT 27892

Score = 930 (139.5 bits), Expect = 2.8e-90, Sum P(2) = 2.8e-90
Identities = 270/354 (76%), Positives = 270/354 (76%), Strand = Minus / Plus

75 Query: 357 TGAAACATCACTGTCT-TGCCAGAGTACCAACACTGTCATGTGATTGATGCCGCCCTCTC 299
|| | | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27795 TGCCAGATTACCGACACTGTCACTTGACCAATACTGACCC-TCTTT-ACCTCGCCACGC 27852

80 Query: 298 TAGAC-CTCACCAC-GCGGACACAT-GCTTCC-G--GTACCTTTGGGTCTGTGAGGTTT 245
||| | | | | | | | | | | | | | | | | | | | | |
Sbjct: 27853 G-GACACCGCTCCT GT|GTCGCTTTGCCCTATTGATGTTCTCTTTGGGTCTGTGAGGTTT 27911

85 Query: 244 TGTCAA--GCGCTAGTGCTAACGCCGTTCTGTACAACCTAACTCACTGGCAAGAACACAG 187
TGT AA G G | TAGTG | T A | G GTT | TGT A | AA | TAA | T | A | TGG | AGAA A | AG
Sbjct: 27912 TGTAAACTGTGCTAGTGCTGACGATGTTCTGTACAACCTAACTCACTGGCGAGAATACAG 27971

90 Query: 186 TGTGTGGGCTTTTCGACCCTAGAACAACTTTTTCAATTGACAGTTGCAGAATTGTGGA 127
GT GG | | TT | | | A | TA AA | A A TTTTT AATTGA | AGTTG | AGAATTGTGGA
Sbjct: 27972 CGTGGGACCTTCAGCCACTACAACAGAATTTTTAAATTGACAGTTGCAGAATTGTGGA 28031

Query: 126 GTGTTTTTACATTGATCTTTTGCTAATGCAGTTAGCAGTATGTTTGCATGTATGACTTA 67
 GTGTTTTTA|ATTGAT|TTTGT|TAATG|A TTAG|A TATGTTTG|ATGTATGA|TTA
 Sbjet: 28032 GTGTTTTTACATTGATCTTTTGCTAATGCAATTAGCATTATGTTTGCATGTATGACTTA 28091
 Query: 66 ATAAATCCTTGAATCATAAAAAAAAAAAAAAAAATGTCTTTGGAACCTGAAAAAAAA 10
 ATAAAT||TTGAAT|ATA A AA A TGT TTTG A|TTGA AA AA
 Sbjet: 28092 ATAAATCCTTGAATCATACGACTGGTAATACTGGTGTTTTGAGACTTGATGAACAA 28148

10 RISKMARKER2

RISKMARKER2 is a 650 bp rat expressed sequence tag (EST) [AW435096]. The nucleic acid sequence was initially identified in a cloned fragment having the following sequence:

```

1 TTTTTTTTTTTTTTTTTTGGCAGAATTCTGATGTTTACTGGGACCCATAGTAGTCAAGGTGACAGCAAGGGTAGGGGAGGA
81 AACTCAGCAGAGGCGGATCCCAGGTCTGGAGGGAAGCTGACAGCAGCCCAGTAAGCTGTGCCAGAAGGCTGTAACAGTAG
161 CGGAGCCAGTGACAGCGCCAGGCTGGGCTGGGTTCTCTGTGGGTGTGCACGGCAAAGCTGCGGCCTGTGGGCCCTGGG
241 GGGCCTGTGACGTCCACATCCACCACATGCATGTCGGTGAGGCTAAGGTGAGCCACAAGCAGCCCCAATGACACGATCAAA
321 GCCTAGACTGGGAGCGGCCAGGGCAGCGGCTGCCATGGTGTGGAGTTTCGGGGGGCCAAGGGGCAGAGCCCACGCACAG
401 GGGCCTCATAGAGCACTGTGCGGGGCCCACTACTATGTGCGGCAGCCAGGGGTCCCTCCAGCCGGAAGCCATCAGGATGT
481 GTGG (SEQ ID NO: 3)

```

15 The cloned sequence was assembled into a contig resulting in the following consensus sequence:

```

1 TTTTTTTTTTTTTTTTTTGGCAGAATTCTGATGTTTACTGGGACCCATAGTAGTCAAGGTGACAGCAAGGGTAGGGGAGGA
81 AACTCAGCAGAGGCGGATCCCAGGTCTGGAGGGAAGCTGACAGCAGCCCAGTAAGCTGTGCCAGAAGGCTGTAACAGTAG
161 CGGAGCCAGTGACAGCGCCAGGCTGGGCTGGGTTCTCTGTGGGTGTGCACGGCAAAGCTGCGGCCTGTGGGCCCTGGG
241 GGGCCTGTGACGTCCACATCCACCACATGCATGTCGGTGAGGCTAAGGTGAGCCACAAGCAGCCCCAATGACACGATCAAA
321 GCCTAGACTGGGAGCGGCCAGGGCAGCGGCTGCCATGGTGTGGAGTTTCGGGGGGCCAAGGGGCAGAGCCCACGCACAG
401 GGGCCTCATAGAGCACTGTGCGGGGCCCACTACTATGTGCGGCAGCCAGGGGTCCCTCCAGCCGGAAGCCATCAGGATGT
481 GTGGCCATGGTGACTCGAAGGCTCTGGAGGCCTCCGGCTGCATCCAATCTGCTGATGTCTTCACAACCCACAGGGCCCC
561 TCGGGCCACAAACACCGTGTGGCCCCAGTGGTTTGAAGCCTCCAGGAGTGCCGCTCTGTGGTCTGGTCAGCGAGAGCTG
641 AGGGGGATCC (SEQ ID NO: 4)

```

Blast-N Results:

20 >gb:GENBANK-ID:AW435096|acc:AW435096 UI-R-BJ0p-afy-e-10-0-UI.s1 UI-R-BJ0p
 Rattus norvegicus cDNA clone UI-R-BJ0p-afy-e-10-0-UI 3', mRNA sequence -
 Rattus norvegicus, 484 bp (RNA).
 Length = 484

25 Plus Strand HSPs:

Score = 2413 (362.0 bits), Expect = 1.2e-102, P = 1.2e-102
 Identities = 483/484 (99%), Positives = 483/484 (99%), Strand = Plus / Plus

30 Query: 1 TTTTTTTTTTTTTTTTTTGGCAGAATTCTGATGTTTACTGGGACCCATAGTAGTCAAGGTG 60
 ||||||||||||||||||||||||||||||||||||||||||||||||||||||||||||
 Sbjet: 1 TTTTTTTTTTTTTTTTTTGGCAGAATTCTGATGTTTACTGGGACCCATAGTAGTCAAGGTG 60

Query: 61 ACAGCAAGGGTAGGGGAGGAACTCAGCAGAGGCGGATCCCAGGTCTGGAGGGAAGCTGA 120
 |||
 5 Sbjct: 61 ACAGCAAGGGTAGGGGAGGAACTCAGCAGAGGCGGATCCCAGGTCTGGAGGGAAGCTGA 120

Query: 121 CAGCAGCCCAGTAAGCTGTGCCAGAAGGCTGTAACAGTAGCGGAGCCAGTGACAGCGCCA 180
 |||
 Sbjct: 121 CAGCAGCCCAGTAAGCTGTGCCAGAAGGCTGTAACAGTAGCGGAGCCAGTGACAGCGCCA 180

10 Query: 181 GGCTGGGCTGGGTTCTCTGTGGGTGTGCACGGCAAAGCTGCGGCCTGTGGGCCCTGGG 240
 |||
 Sbjct: 181 GGCTGGGCTGGGTTCTCTGTGGGTGTGCACGGCAAAGCTGCGGCCTGTGGGCCCTGGG 240

15 Query: 241 GGGCCTGTCTAGTCCACATCCACCACATGCATGTCTGGTGAGGCTAAGGTCAGCCACAAGC 300
 |||
 Sbjct: 241 GGGCCTGTCTAGTCCACATCCACCACATGCATGTCTGGTGAGGCTAAGGTCAGCCACAAGC 300

20 Query: 301 ACCCAATGACACGATCAAAGCCTAGACTGGGAGCGGCCAGGGCAGCGGCTGCCATGGTG 360
 |||
 Sbjct: 301 ACCCAATGACACGATCAAAGCCTAGACTGGGAGCGGCCAGGGCAGCGGCTGCCATGGTG 360

25 Query: 361 TTGGAGTTTCGGGGGGCCAAGGGGAGAGCCACGCACAGGGCCCTCATAGAGCACTGTG 420
 |||
 Sbjct: 361 TTGGAGTTTCGGGGGGCCAAGGGGAGAGCCACGCACAGGGCCCTCATAGAGCACTGTG 420

30 Query: 421 CGGGGGCCCACTACTATGTGCGGCAGCCAGGGGTCCCTCCAGCCGGAAGCCATCAGGATGT 480
 |||
 Sbjct: 421 CGGGGGCCCACTACTATGTGCGGCAGCCAGGGGTCCCTCCAGCCGGAAGCCATCAGGATGT 480

30 Query: 481 GTGG 484
 |||
 Sbjct: 481 GTGG 484

35 Blast-X Results:

>ptnr:SPTREMBL-ACC:Q19527 F17C8.3 PROTEIN - *Caenorhabditis elegans*, 973 aa.

[Top](#) [Previous Match](#) [Next Match](#)
 Length = 973

Minus Strand HSPs:

40

Score = 351 (123.6 bits), Expect = 6.3e-30, P = 6.3e-30
 Identities = 78/161 (48%), Positives = 96/161 (59%), Frame = -1

45 Query: 650 GSPSALADQTTERRQLLEASNHWGHTVFVARGALWGCEISRDLAAGGLQSLRVMTATHPD 471
 GSP+ A+Q +L+S G ++ GALWG DI++ GL+LVTM HP
 Sbjct: 530 GSPTCFANQELLEKLTKLSLSHGKLLIPAGALWGANDIQKMADVGLKGLTVMIKHPT 589

50 Query: 470 GFRLEGPLAAAHSSGP-----RTVLYEGPVRGLCPLAPRNSNTMAAALAAPSLGFDRVI 306
 F+L PL + TVLYEG VRGLCPLAP N NTMA ALAA +LGFD V
 Sbjct: 590 SFKLGSPLFEINEKAKLEETNETVLYEGSVRGLCPLAPNNVNTMAGGALAASNLFDEVK 649

55 Query: 305 GVLVADLSLTDHVVVDVELTGPPGPTGRSFAVHTHRENPAPGAVT 168
 L++D +TD HVV+V+G G F V T R NPA+PGAVT
 Sbjct: 650 AKLISDPKMTDWHVVEVRVEGDDG-----FEVITRRNNPAKPGAVT 690

RISKMARKER3

RISKMARKER3 is a 1019 nucleotide sequence encoding superoxide dismutase copper chaperone [AF255305]:

1 ggtctctgga ccctaccggt tgtgtggccc aagcgggtga ctgcagccag gatggcttcg
 60 61 aagtcggggg acggtggaac tatgtgtgcg ttggagttaa cagtacagat gagttgtcag
 121 agctgcgtgg acgctgtgca caagaccctg aaaggggagg cggtgttcca gaatgtggaa
 181 gttcagttgg agaaccagat ggtgttggtg cagaccactt tgcccagcca ggaggtgcaa
 241 gcgctcctgg aaagcacagg gaggcaggct gtactcaagg gcatgggcag cagccaacta
 301 aagaatctgg gagcagcagt ggccattatg gagggcagtg gcaccgtaca gggggtggtc
 65 361 cgcttcttac agctgtcctc tgagctctgc ctgattgagg gaaccatcga cggcctggag
 421 cctgggctgc atgggcttca tgtccatcag tatggggacc ttaccaagga ctgcagcagc

```

481  tgtggggacc attttaaccc tgatggagca tctcatgggg gtcctcagga
cactgatcgg 541  caccggggag atctgggcaa tggtcacgct gaagctagt
gccgagctac cttccggata 601  gaggataaac agctgaaggt gtgggatgtg
attggccgca gtctggtgtg tgatgaggga 661  gaagatgacc tgggcccggg
5  aggccatccc ttatccaagg tcacaggga ttctgggaag 721  aggttggcct
gtggcatcat tgcacgctct gctggccttt tccagaatcc caagcagatc 781  tgctcctgtg
atgggctcac tatctgggag gaggcaggcc ggccattgc tggccaaggc 841  cgaaaggact
cagcccaacc ccctgctcac ctctgaacag agcctcctgt caggttattc 901  agtcctccta
gctgaacatc ttcctgcaga gggagcctca agcccttgct tgtataggcc 961  taaagggcag
10 ataggcattg ttgtatcctg agcaaattaa attgttactc tcatatggc

```

RISKMARKER4

RISKMARKER4 is a 878 nucleotide sequence encoding alpha-2 microglubulin [U31287]:

```

15  1  ggcacgagca gagagattgt cccaacagag aggcaattct attccctacc
aacatgaagc
61  tgttgctgct gctgctgtgt ctgggcctga cactggctctg tggccatgca
gaagaagcta 121  gttccacaag agggaacctc gatgtggcta agctcaatgg
ggattggttt tctattgtcg 181  tggcctctaa caaaagagaa aagatagaag
20  agaatggcag catgagagtt tttatgcagc 241  acatcgatgt cttggagaat
tccttaggct tcaagttccg tattaaggaa aatggagagt 301  gcagggaact
atatttggtt gcctacaaaa cgccagagga tggcgaatat tttgttgagt 361  atgacggagg
gaatacattt actatactta agacagacta tgacagatat gtcattgttc 421  atctcattaa
tttcaagaac ggggaaacct tccagctgat ggtgctctac ggcagaacaa 481  aggatctgag
25  ttcagacatc aaggaaaagt ttgcaaaact atgtgaggcg catggaatca 541  ctagggacaa
tatcattgat ctaaccaaga ctgatcgctg tctccaggcc cgaggatgaa 601  gaaaggcctg
agcctccagt gctgagtgga gacttctcac caggactcta gcatcaccat 661  ttcctgtcca
tgagacatcc tgagacaaat tctgcatctc gatttccatc ctctgtcaca 721  gaaaagtcca
atcctggctc ctccagcatc ttccctaggt taccaggac aacacatcga 781  gaattaaaag
30  ctttcttaaa tttctcttgg cccaccccat gatcattccg cacaatatc 841  ttgctcttgc
agttcaataa atgattaccc ttgcactt

```

RISKMARKER5

RISKMARKER5 is a 2443 bp rat mRNA for Mx3 protein [X52713]. The nucleic acid was initially identified in a cloned fragment (having 100% sequence identity to the rat mRNA) having the following sequence:

```

1  CCATGGATGAAATCTTCCAGCATCTGAATGCCTACCGCCAGGAGGCTCACAACTGCATCTCCAGCCACATTCCATTGATC
81  ATCCAGTATTTTCATCTTGAAGATGTTTGCTGAGAAGCTGCAGAAGGGCATGCTCCAGCTCCTGCAGGACAAGGATTCCTG
161  CAGCTGGCTCCTGAAGGAAAAGAGTGACACCACTGAGAAGAGGAGATTCTCTGAAGGAGCGGTTGGCAAGGCTGGCCCAAG
241  CTCAGCGCAGGCTAGC (SEQ ID NO: 5)

```

Blast-N Results:

```

>gb:GENBANK-ID:RNMX3|acc:X52713 Rat mRNA for Mx3 protein - Rattus
norvegicus, 2443 bp.
Top Previous Match Next Match
Length = 2443

```

Plus Strand HSPs:

Score = 1280 (192.1 bits), Expect = 9.5e-52, P = 9.5e-52
Identities = 256/256 (100%), Positives = 256/256 (100%), Strand = Plus / Plus

5 Query: 1 CCATGGATGAAATCTTCCAGCATCTGAATGCCTACCGCCAGGAGGCTCACAACCTGCATCT 60
|||||
Sbjct: 1710 CCATGGATGAAATCTTCCAGCATCTGAATGCCTACCGCCAGGAGGCTCACAACCTGCATCT 1769

10 Query: 61 CCAGCCACATTCCATTGATCATCCAGTATTTTCATCTTGAAGATGTTTGCTGAGAAGCTGC 120
|||||
Sbjct: 1770 CCAGCCACATTCCATTGATCATCCAGTATTTTCATCTTGAAGATGTTTGCTGAGAAGCTGC 1829

15 Query: 121 AGAAGGGCATGCTCCAGCTCCTGCAGGACAAGGATTCCTGCAGCTGGCTCCTGAAGGAAA 180
|||||
Sbjct: 1830 AGAAGGGCATGCTCCAGCTCCTGCAGGACAAGGATTCCTGCAGCTGGCTCCTGAAGGAAA 1889

20 Query: 181 AGAGTGACACCAGTGAGAAGAGGAGATTCTGAAGGAGCGGTTGGCAAGGCTGGCCCAAG 240
|||||
Sbjct: 1890 AGAGTGACACCAGTGAGAAGAGGAGATTCTGAAGGAGCGGTTGGCAAGGCTGGCCCAAG 1949

25 Query: 241 CTCAGCGCAGGCTAGC 256
|||||
Sbjct: 1950 CTCAGCGCAGGCTAGC 1965

Blast-X Results:

>ptnr:SWISSPROT-ACC:P18590 INTERFERON-INDUCED GTP-BINDING PROTEIN MX3 -
Rattus norvegicus (Rat), 659 aa.

Top Previous Match Next Match
Length = 659

30 Plus Strand HSPs:

Score = 429 (151.0 bits), Expect = 5.3e-39, P = 5.3e-39
Identities = 84/84 (100%), Positives = 84/84 (100%), Frame = +3

35 Query: 3 MDEIFQHLNAYRQEAHNCISSHIPLIIQYFILKMFAEKLQKGMQLQLQDKDSCSWLLKEK 182
MDEIFQHLNAYRQEAHNCISSHIPLIIQYFILKMFAEKLQKGMQLQLQDKDSCSWLLKEK
Sbjct: 571 MDEIFQHLNAYRQEAHNCISSHIPLIIQYFILKMFAEKLQKGMQLQLQDKDSCSWLLKEK 630

40 Query: 183 SDTSEKRRFLKERLARLAQAQRRL 254
SDTSEKRRFLKERLARLAQAQRRL
Sbjct: 631 SDTSEKRRFLKERLARLAQAQRRL 654

RISKMARKER6

45 RISKMARKER6 is 369 bp novel gene fragment, which has 98% amino acid identity (90% nucleic acid sequence identity) to Human ERj3 protein [AJ250137]. The nucleic acid sequence was initially identified in a cloned fragment having the following sequence:

1 TCTAGAAAGTCACCTTGGGAAGAAGTGACGCAGGGAACCTTTGTGGAAGTAGTTAGAAACAAGCCCGTAGCCAGGCAGGCT
81 CCTGGCAAACGGAAATGCAACTGTCTGGCAGGAGATGCGAACCACACAGCTGGGACCAGGGCGCTTCCAAATGACCCAGGA
161 AGTGGTTTGTGACGAGTGCCCTAATGTCAAAGTAGTGAATGAAGAACGAACACTAGAAAGTGGAATAGAGCCTGGGGTGA
241 GAGATGGCATGGAGTACCCCTTTATTGGAGAAGGTGAGCCTCATGTGGATGGGGAACCCGGAGACTTACGGTTCCGAATC
321 AAAGTTGTCAAGCACCAGGATATTTGAGAGGAGAGGGGATGACCTGTACA (SEQ ID NO: 6)

Blast-N Results:

>gb:GENBANK-ID:HSA250137|acc:AJ250137 Homo sapiens mRNA for ERj3 protein (ERj3 gene) - Homo sapiens, 1159 bp.

50 Top Previous Match Next Match
Length = 1159

Plus Strand HSPs:

55 Score = 1524 (228.7 bits), Expect = 5.6e-63, P = 5.6e-63

Identities = 334/369 (90%), Positives = 334/369 (90%), Strand = Plus / Plus

Query: 1 TCTAGAAAGTCACCTTGGGAAGAAGTGTACGCAGGGAACCTTTGTGGAAGTAGTTAGAAACA 60
 TCTAGAA GTCAC TTGGAAGAAGT TA GCAGG AA TTTGTGGAAGTAGTTAGAAACA
 5 Sbjct: 431 TCTAGAA-GTCACCTTGGGAAGAAGTATATGCAGGAAATTTGTGGAAGTAGTTAGAAACA 489

Query: 61 AGCCCGTAGCCAGGCAGGCTCCTGGCAAACGGAATGCAACTGTGCGCAGGAGATGCGAA 120
 A CC GT GC AGGCAGGCTCCTGGCAAACGGA TGCAA TGTCGGCA GAGATGCG A
 10 Sbjct: 490 AACCTGTGGCAAGGCAGGCTCCTGGCAAACGGAAGTGCATGTGCGCAAGAGATGCGGA 549

Query: 121 CCACACAGCTGGGACCAGGCGCTTCCAAATGACCCAGGAAGTGGTTTGTGACGAGTGCC 180
 CCAC CAGCTGGG CC GGGCGCTTCCAAATGACCCAGGA GTGGT TG GACGA TGCC
 Sbjct: 550 CCACCCAGCTGGGCCCTGGGCGCTTCCAAATGACCCAGGAGGTGGTCTGCGACGAATGCC 609

Query: 181 CTAATGTCAAACCTAGTGAATGAAGAACGAACACTAGAAAGTGGAAATAGAGCCTGGGGTGA 240
 CTAATGTCAAACCTAGTGAATGAAGAACGAAC CT GAAGT GAAATAGAGCCTGGGGTGA
 15 Sbjct: 610 CTAATGTCAAACCTAGTGAATGAAGAACGAACGCTGGAAGTAGAAATAGAGCCTGGGGTGA 669

Query: 241 GAGATGGCATGGAGTACCCCTTTATTGGAGAAGGTGAGCCTCATGTGGATGGGGAACCCG 300
 GAGA GGCATGGAGTACCCCTTTATTGGAGAAGGTGAGCCTCA GTGGATGGGA CC G
 20 Sbjct: 670 GAGACGGCATGGAGTACCCCTTTATTGGAGAAGGTGAGCCTCACGTGGATGGGAGCCTG 729

Query: 301 GAGACTTACGGTTCCGAATCAAAGTTGTCAAGCACCGATATTTGAGAGGAGAGGGGATG 360
 GAGA TTACGGTTCCGAATCAAAGTTGTCAAGCACC ATATTTGA AGGAGAGG GATG
 25 Sbjct: 730 GAGATTTACGGTTCCGAATCAAAGTTGTCAAGCACCAATATTTGAAAGGAGAGGAGATG 789

Query: 361 ACCTGTACA 369
 A TGTACA
 Sbjct: 790 ATTTGTACA 798
 30

Blast-X Results:

>ptnr:SPTREMBL-ACC:Q9UBS4 ERJ3 PROTEIN PRECURSOR - Homo sapiens
 (Human), 358 aa.

Top Previous Match Next Match
 Length = 358

35 Plus Strand HSPs:

Score = 637 (224.2 bits), Expect = 2.1e-61, P = 2.1e-61
 Identities = 119/121 (98%), Positives = 120/121 (99%), Frame = +3

40 Query: 6 KVTLEEVYAGNFVEVVRNKPVARQAPGKRKCNCRQEMRTTQLGPGRFQMTQEVVCDCEPN 185
 +VTLEEVYAGNFVEVVRNKPVARQAPGKRKCNCRQEMRTTQLGPGRFQMTQEVVCDCEPN
 Sbjct: 139 EVTLEEVYAGNFVEVVRNKPVARQAPGKRKCNCRQEMRTTQLGPGRFQMTQEVVCDCEPN 198

45 Query: 186 VKLVNEERTLEVEIEPGVRDGMIEYFFIGEGEPHVDGEPGDLRFRIKVVKHIFERRGDDL 365
 VKLVNEERTLEVEIEPGVRDGMIEYFFIGEGEPHVDGEPGDLRFRIKVVKH IFERRGDDL
 Sbjct: 199 VKLVNEERTLEVEIEPGVRDGMIEYFFIGEGEPHVDGEPGDLRFRIKVVKHIFERRGDDL 258

Query: 366 Y 368
 Y
 50 Sbjct: 259 Y 259

RISKMARKER7

RISKMARKER7 is a 594 bp novel gene fragment, which has 65% sequence identity to
Mus musculus hexokinase II [AJ238540], probable 3' UTR. The nucleic acid sequence was
 55 initially identified in a cloned fragment having the following sequence:

1 GGGCCCCACTAAACATACACAAAAGAATAAAATGTTTCATTTAACTTAACTGCTTCCTGGTTTTACAAGGCATAAA
 81 TATATAGCATCTCCAACAGCTACCTGTAGATTCTGTTAGTGCAAAACCTTAGAAACCTCCTGGAGCTCAAAGGCATCCG
 161 GACTAGT (SEQ ID NO: 7)

The cloned sequence was assembled into a contig resulting in the following consensus
 sequence:

1 TTTTTTTTTTTTTTTTAAAAAGATTATAAAATTGAATTTATTGAGTTTCACACAAGATGCACCTATAAAATTAGTACT
81 GAATGCCATTATGACAGAAGTGAGCATCATCCACACTCCCAAGAGCATCTGCAAAGGAAATCAATCTTCAGAGAATAGCA
161 CAGAAACAGAAAATCCAAGCGAACAAAAAGATACATCTAGGCCGTGTTCTTGTCTGACCAGGGCCGCATTTGGCAAAGC
241 TTTCTCTGCACCTCCCCTGGTTGCCAAGGATACTTTCTTTTGTAAAAAAAAGTTTAGAAGTGGGGCCCCACTAAAC
321 ATACACAAAAGAATAAAAATGTTCACTTTAACTTAACTGCTTCCTGGTTTACAAGGCATAAATATATAGCATCTCCA
401 ACAGCTACCTGTAGATTCTGTGTAGTGCAAAACCTTAGAAACCTCCTGGAGCTCAAAGGCATCCGGACTAGTTTTGTACT
481 TAAACAGGATACGGGTAAACCACTTAAATTTGCCATCTCTGCCCAAAGTGTTCATGAGAACTGAGTTTCAGAAGACA
561 GCATAGGAAAGAGTCAGAAACGGTCACTTTTTT (SEQ ID NO: 8)

Blast-N Results:

>gb:GENBANK-ID:MMU238540|acc:AJ238540 Mus musculus mRNA for hexokinase
II - Mus musculus, 5474 bp.

Top Previous Match Next Match
Length = 5474

5 Minus Strand HSPs:

Score = 251 (37.7 bits), Expect = 0.045, P = 0.044
Identities = 121/184 (65%), Positives = 121/184 (65%), Strand = Minus / Plus

10 Query: 184 GTTCGCTTGGATTTT-CTG-TTCTGTGCTATTCTCTGAAGATT-GATTTCCTTTGCAGA 128
G TC CT G T T CTG TTT TGTG T TTC TGAA TT GA T C T T CA A
Sbjct: 5287 GCTCTCTCTGCTAATGCTGCTTTGTGTGATCTTCAGTGAACCTTTGACT-CATCT-CATA 5344

15 Query: 127 TGCTCTTGGGAGTGTGGATGATGCTCACTTCTGTCATAATGG-CATTC-AGTACTAATTT 70
T C CT GG A T G T TG C TT TGTCAT ATG CA T AG ACTA TT
Sbjct: 5345 TCC-CTGGGCACCTCGGTCTAGTGAGCGTTT-TGTCATCATGTACAGTAGAGAACTAGTTG 5402

20 Query: 69 TATAAGTGCATCTTGTGTGAACTCA-ATAAATCAATTTTATAATCTTTTTTAAAAAAA 11
AT A CAT T TGT AA CT AT AAT AATTTTA T TTTTTT AAAAAA
Sbjct: 5403 AATTAAC-CATGTGATGTTAA-CTATTATTAATA-AATTTTAACCTTTTTTTTCAAAAAA 5459

Query: 10 AAAAAAAAAA 1
AAAAAAAAAA
Sbjct: 5460 AAAAAAAAAA 5469

25 Score = 250 (37.5 bits), Expect = 0.051, P = 0.049
Identities = 122/184 (66%), Positives = 122/184 (66%), Strand = Minus / Plus

30 Query: 184 GTTCGCTTGGATTTT-CTG-TTCTGTGCTATTCTCTGAAGATT-GATTTCCTTTGCAGA 128
G TC CT G T T CTG TTT TGTG T TTC TGAA TT GA T C T T CA A
Sbjct: 5287 GCTCTCTCTGCTAATGCTGCTTTGTGTGATCTTCAGTGAACCTTTGACT-CATCT-CATA 5344

35 Query: 127 TGCTCTTGGGAGTGTGGATGATGCTCACTTCTGTCATAATGG-CATTC-AGTACTAATTT 70
T C CT GG A T G T TG C TT TGTCAT ATG CA T AG ACTA TT
Sbjct: 5345 TCC-CTGGGCACCTCGGTCTAGTGAGCGTTT-TGTCATCATGTACAGTAGAGAACTAGTTG 5402

40 Query: 69 TATAAGTGCATCTTGTGTGAACTCA-ATAAATCAATTTTATAATCTTTTTT-AAAAAA 12
AT A CAT T TGT AA CT AT AAT AATTTTA T TTTTTT AAAAAA
Sbjct: 5403 AATTAAC-CATGTGATGTTAA-CTATTATTAATA-AATTTTAACCTTTTTTTTCAAAAAA 5459

Query: 11 AAAAAAAAAA 1
AAAAAAAAAA
Sbjct: 5460 AAAAAAAAAA 5470

45 Blast-X Results:

>ptnr:SPTREMBL-ACC:Q9VIA2 MST84DB PROTEIN - Drosophila melanogaster
(Fruit fly), 70 aa.

Top Previous Match Next Match
Length = 70

50 Plus Strand HSPs:

Score = 66 (23.2 bits), Expect = 2.2, P = 0.88
Identities = 15/48 (31%), Positives = 25/48 (52%), Frame = +3

Query: 66 YKISTECHYDRSEHHPHSQEHLQRKS-----IFRE*HNRKSKRTKR 191
 YK+ ++ H R +H P S++ RK I ++ RNRK R ++
 Sbjct: 3 YKVHSHVHKARMDHSPRSKDRKDRKGRKAHSKIHKDYSRNRKDHVRK 50

5

RISKMARKER8

RISKMARKER8 is a 797 bp novel gene fragment, which has 94% amino acid identity (79% nucleic acid sequence identity) to human GT335 mRNA (ES1 Protein Homolog) [U53003]. The nucleic acid was initially identified in a cloned fragment having the following sequence:

10

```

1 CCTAGGACTGCACAACGTGAGTCCTTGAACCAGGCTCTGGAAGGTGCCAGACCACCAATGGGGACACAGTGAGG
81 CCAGCCCCCAGTGAAATTCCTGCTGCTACCTGGGGCCCTTGGTGAGACTCTGGCTTCCGGCTGGTAGAAGCCAAGGTTGG
161 ACGCATAGTTGCAAAGCTCCTCCTTCAGGCACAAAGTGCTATGCTTCTAATAGAACAGCAGCTCCCGTGTCTGGCTGA
241 CCGGAGCACACAGGCTGAGCGTGCCACAGCGACGAGGCCAAGCGTGGTGCTGGTGGTGTACTTTCCCGTGAGTTCC
321 CAGCACCTTCTTCACCATGG (SEQ ID NO: 9)

```

The cloned sequence was assembled into a contig resulting in the following consensus sequence:

```

1 TTTTTTTTTTTTTTTTTTTTTTGTAGTTTCCACTGTGGAAGAGTTTATTGTATGGCTGCAGGGATCTACTACAGAATCC
81 CCCTGGCTGCAGTTAGCTGTGCTTACTCTGGACATATCTCCGAAGACTTGGAGCCTAAATGGTTTCTTCTTTAGAGCT
161 TTAGTACCCGATCCATCAGACCTAGGACTGCACAACGTGAGTCCTTGAACCAGGCTCTGGAAGGTGCCAGACCACCC
241 AATGGGGACACACAGTGAGGCCAGCCCCAGTGAAATTCCTGCTGCTACCTGGGGCCCTTGGTGAGACTCTGGCTTCCGG
321 CTGGTAGAAGCCAAGGTTGGACGCATAGTTGCAAAGCTCCTCCTTCAGGCACAAAGTGCTATGCTTCTAATAGAACAGC
401 AGCTCCCGTGTCTGGCTGACCGGAGCACACAGGCTGAGCGTGCCACAGCGACGAGGCCAAGCGTGGTGCTGGTGG
481 TGTTACTTTCCCGTGAGTTCCAGCACCTTCTTCACCATGGCCCCAATCCCGTCGTGGATGTGGTGGAGTTCGGTCTCACA
561 CATGAAGGCCGGGGTGGTGACCACCTTGTCTTCTGGTCGACGTGAGCTTCGGTCACACCCTTCACACAGTGCTTGGCAC
641 CCAGGGCTTTGACGGCTTCCGCGTTCCAGCATATGGCCACTTGCCCCCTCCTTTGCTCATGGCCACGGTGACCTCC
ACACCTTTGATCACTTTGGCTGCGAGGACAGGAGCGATGCAGCATAGGCCAATGGGCTTCTTGGCTCCGTGGAATTC
721 (SEQ ID NO: 10)

```

Blast-N Results:

>gb:GENBANK-ID:HSU53003|acc:U53003 Human GT335 mRNA, complete cds - Homo sapiens, 1652 bp.

15

Top Previous Match Next Match
 Length = 1652

Minus Strand HSPs:

20

Score = 1141 (171.2 bits), Expect = 7.9e-46, P = 7.9e-46
 Identities = 307/385 (79%), Positives = 307/385 (79%), Strand = Minus / Plus

Query: 797 GAATTCACGGAGCCAAGAAGCCCATTTGGCCTATGCTGCATCGCTCCTGCTCGCAGCC 738
 GA TTCCAC GCC GAAGCCCAT GGC T TGCTGCAT GC CTTGCTCGC GCC
 Sbjct: 577 GAGTTCCACAGGCCGGGAAGCCCATCGGCTTGTGCTGCATTGCACCTGCTCCTCGCGGCC 636

25

Query: 737 AAAGTGATCAAAGGTGTGGAGGTCACCGTGGGCCATGAGCAAGAGGAGGGGGCAAGTGG 678
 AA GTG TCA AGG GT GAGGT AC GTGGGCCA GAGCA GAGGA GG GGCAAGTGG
 Sbjct: 637 AAGGTGCTCAGAGGCGTCGAGGTGACTGTGGGCCACGAGCAGGAGGAAGGTGGCAAGTGG 696

5 Query: 677 CCATATGCTGGAACCGCGGAAGCCGTCAAAGCCCTGGGTGCCAAGCACTGTGTGAAGGGT 618
 CC TATGC GG ACCGC GA GCC TCAA GCCCTGGGTGCCAAGCACTG GTGAAGG
 Sbjct: 697 CCTTATGCCGGGACCGCAGAGGCCATCAAGGCCCTGGGTGCCAAGCACTGCGTGAAGGAA 756

10 Query: 617 GTGACCGAAGCTCACGTCGACCAGAAAAACAAGGTGGTCAACACCCCGCCTTCATGTGT 558
 GTG CGAAGCTCACGT GACCAGAAAAACAAGGTGGTCAAC ACCCC GCCTTCATGTG
 Sbjct: 757 GTGGTCGAAGCTCACGTGGACCAGAAAAACAAGGTGGTCAACACCCCGCCTTCATGTGC 816

15 Query: 557 GAGACCGAAGCTCACACCATCCACGACGGGATTGGGGCCATGGTGAAGAAGGTGCTGGAA 498
 GAGAC G ACTCCAC ACATCCA GA GGGAT GG GCCATGGTGA GAAGGTGCTGGAA
 Sbjct: 817 GAGACGGCACTCCACTACATCCATGATGGGATCGGAGCCATGGTGAAGAAGGTGCTGGAA 876

20 Query: 497 CTCACGGGAAAGTAACAC-CACC-A-GCACCAC-GCTTGGCCTCCGT-CGTCGCTGTGGC 443
 CTCAC GGAAAGT AC C CA A G C C GCT GGC C G C T GC T C
 Sbjct: 877 CTCACTGGAAGTGACGCGCATGGACGGGGCCAGCTAGGCGCCAGGACTTGGCC-T--C 933

25 Query: 442 ACGCTCAGCCTGTGT-GCTC-CGGTCAGC 416
 AC CTC G CTG G GCT CGG C GC
 Sbjct: 934 ACCCTCTGGCTGAGGAGCTGTCGG-CTGC 961

25 **Blast-X Results:**
 >ptnr:SWISSNEW-ACC:P30042 ES1 PROTEIN HOMOLOG, MITOCHONDRIAL PRECURSOR
 (PROTEIN KNP-I) (GT335 PROTEIN) - Homo sapiens (Human), 268 aa.
 Top Previous Match Next Match
 Length = 268

30 Minus Strand HSPs:
 Score = 505 (177.8 bits), Expect = 2.0e-47, P = 2.0e-47
 Identities = 94/104 (90%), Positives = 99/104 (95%), Frame = -1

35 Query: 797 EFHGAKKPIGLCCIAPVLAAKVIKGVETVVGHEQEEGGKWPYAGTAEAVKALGAKHCVK 618
 EFH A KPIGLCCIAPVLAAKV++GVEVTVGHEQEEGGKWPYAGTAEA+KALGAKHCVK
 Sbjct: 165 EFHQAGKPIGLCCIAPVLAAKVLRGVETVVGHEQEEGGKWPYAGTAEAIKALGAKHCVKE 224

40 Query: 617 VTEAHVDQKNKVVTTPAFMCETELHHIHDGIGAMVKVLELTGK 486
 V EAHVDQKNKVVTTPAFMCET LH+IHDGIGAMV+KVLELTGK
 Sbjct: 225 VVEAHVDQKNKVVTTPAFMCETALHYIHDGIGAMVRKVLELTGK 268

Principle components analysis was used to generate three eigenvectors used to transform the original expression level data matrix, as shown in Table 4 below. Eigenvector 1 values represent NSAIDs with low risk of hepatotoxicity, Eigenvector 2 values represent NSAIDs with very low risk of hepatotoxicity, and Eigenvector 3 values represent NSAIDs with overdose risk of hepatotoxicity.

Table 4: Transform Eigenvectors for Hepatotoxicity Markers by Risk Classification

Gene	Eigenvector 1	Eigenvector 2	Eigenvector3
RISKMARKER1	26.9	6.7	-0.9
RISKMARKER2	23.3	-1.4	1.5
RISKMARKER3	-26.0	-1.5	-2.3
RISKMARKER4	12.6	-2.2	-6.4
RISKMARKER5	18.0	-1.3	-3.1
RISKMARKER6	-13.8	4.71	19.3
RISKMARKER7	-29.7	-7.5	1.3
RISKMARKER8	19.3	1.2	-2.6
% of variation explained	99.6	0.4	0.1

These eigenvectors may be used to transform the expression levels of RISKMARKERS 1-8 ("RISKMARKERS") in response to a given drug, in order to determine that drug's hepatotoxicity risk. For example, expression levels of RISKMARKERS correlating with Eigenvector 1 indicates that the test drug has a low risk of hepatotoxicity.

- 5 Alternatively, a drug's RISKMARKERS expression profile can be generated simultaneously with the above-described training set (or an equivalent set) run in parallel with the test drug, and expression levels associated with the test drug directly compared to those of the training set.

10 A second training set based on type of injury (hepatocellular damage, cholestasis, elevated transaminase level) was also constructed, utilizing the compounds indicated in Table 5, below.

Table 5: Training Set of NSAIDs by Injury Type

Control	Hepatocellular	Cholestasis	Elevated transmainases
Sterile water	Acetaminophen	Benoxaprofen	Zomepirac
10% ethanol	Flurbiprofen	Nabumetone	Mefenamic acid
Canola oil	Ketoprofen	Sulindac	Tenoxicam

This analysis produced ten fragments that significantly ($p=8.7 \times 10^{-18}$) discriminated among the drugs in the test set. The identities of these ten fragments (INJURYMARKER 1-10) that are included in the discriminatory set (with GenBank accession numbers) are shown below. Where appropriate, the cloned sequence from isolation is provided, and this sequence was then extended using either Genbank rat ESTs or from internally sequenced (Curagen Corporation) rat fragments. The fragments were used to extend the cloned sequence, and the extended contig sequence is provided as "consensus." Finally, the best BlastN and BlastX results are also provided. In some instances the cloned sequence is identical to a known rat gene, in those instances the name of the gene and the GenBank accession number is provided.

INJURYMARKER1

INJURYMARKER1 is a 1025 bp rat express sequence tag (EST) [AI169175]. The nucleic acid was initially identified in a cloned fragment having the following sequence:

```

1  CTGATTTCAAATTTTATTAGAGAACACTTTCGGATTTCAAATTTTATTACAGAACAAACATTTCTGATTTCAAATTT
81  CTATTATAATTCTCCAGTAATCAAAGCAGTGGCGTTGGCATGAAGGCAGACAGAGGTCATGGAAGAGACCAGGCTCAGAA
161 ACAGCCCCACCATGCACAGCGGGATGTTTTCCACCAAGGGCAACATGCAAAGCCAGGTATCCACATGGGTAGAGTAGAA

```

241 AGTCAGACCTTACATCTCACACACAAATGAACTCAAATATACCAGAGAGCAAAGCTAAGAGCTAAAATCAAGTTTCCTA
 321 GGGCAAGCTGTAGTAGGCTCCCTTGGGTGGGTTAATGCTTTTGTGGATGTGACTACCAAAAATTCAACCAGAGCCAACGA
 401 CCCAACTATTAATGGGCAGTGGACCTAAAGAGATTCTTCAAACGATATATAAAGAAGGCCACCAAGCATATAAACATG
 481 TGACATCAGTAGTCAGAGAGATGGGAAGCAGAAGCACTAGCAGATCTTAACACCTACTAGAACANCCACTAAAAAAGAGT
 561 AAGACTCACAAGGACATGGGCACTTCTAATCTCTGTGCACTGCTGCCAGGACATACAATAGTGTGGTCACTATGGAGACT
 641 ACGGCAGTGCCTACTAATAACAGCAGAGTTACCCTAAGACATACAATCTGCTGCGTGTATGCTAAGCAGGATCCGAGGGA
 TATTTGTATATACATGTTACAGCATAGTCAGGAGCTCCAGGGTGGGAACAACAGGTACC
 721 (SEQ ID NO: 11)

The cloned sequence was assembled into a contig resulting in the following consensus sequence:

1 CTGATTTCAAATTTTATTATAGAACACTTTCTGATTTCAAATTTTATTACAGAACAAACATTTCTGATTTCAAATTT
 81 CTATTATAATTCTCCAGTAATCAAAGCAGTGGCGTTGGCATGAAGGCAGACAGAGGTCATGGAAGAGACCAGGCTCAGAA
 161 ACAGCCCCACCATGCACAGCGGGATGTTTCCCACCAAGGGCAACATGCAAAGCCAGGTATCCACATGGGTAGAGTAGAA
 241 AGTCAGACCTTACATCTCACACACAAATGAACTCAAATATACCAGAGAGCAAAGCTAAGAGCTAAAATCAAGTTTCCTA
 321 GGGCAAGCTGTAGTAGGCTCCCTTGGGTGGGTTAATGCTTTTGTGGATGTGACTACCAAAAATTCAACCAGAGCCAACGA
 401 CCCAACTATTAATGGGCAGTGGACCTAAAGAGATTCTTCAAACGATATATAAAGAAGGCCACCAAGCATATAAACATG
 481 TGACATCAGTAGTCAGAGAGATGGGAAGCAGAAGCACTAGCAGATCTTAACACCTACTAGAACAGCCACTAAAAAAGAGT
 561 AAGACTCACAAGGACATGGGCACTTCTAATCTCTGTGCACTGCTGCCAGGACATACAATAGTGTGGTCACTATGGAGACT
 641 ACGGCAGTGCCTACTAATAACAGCAGAGTTACCCTAAGACATACAATCTGCTGCGTGTATGCTAAGCAGGATCCGAGGGA
 721 TATTTGTATATACATGTTACAGCATAGTCAGGAGCTCCAGGGTGGGAACAACAGGTACCCACGGCTGGATGAGTAGG
 801 TAACAAGAAACATACAGCATACATACAACACACACTAAAGTCTAAAGTACTATTTGTCCTTACAAAGGAACTCATACAT
 881 GATACAAGCCTTCACGGCATTCTGCTACATGAACACGCACACACACACACACACACACACACACACACGCACTGAGAATC
 TATGTATACCAGGCACCTTAGGGTACTCAAATTCAGAAACAGGACAGAGAATGGTGATTGCCATGG
 961 (SEQ ID NO: 12)

5 Blast-N Results:

>gb:GENBANK-ID:AI169175|acc:AI169175 EST215009 Normalized rat kidney, Bento Soares Rattus sp. CDNA clone RKIBO44 3' end, mRNA sequence - Rattus sp., 670 bp (RNA).

Top Previous Match Next Match
 Length = 670

Plus Strand HSPs:

10

Score = 3305 (495.9 bits), Expect = 4.3e-143, P = 4.3e-143
 Identities = 661/661 (100%), Positives = 661/661 (100%); Strand = Plus / Plus

15

Query: 4 ATTTCAAATTTTATTATAGAACACTTTCTGATTTCAAATTTTATTACAGAACAAACAT 63
 |||
 Sbjct: 1 ATTTCAAATTTTATTATAGAACACTTTCTGATTTCAAATTTTATTACAGAACAAACAT 60

20

Query: 64 TTTCTGATTTCAAATTTCTATTATAATTCTCCAGTAATCAAAGCAGTGGCGTTGGCATGA 123
 |||
 Sbjct: 61 TTTCTGATTTCAAATTTCTATTATAATTCTCCAGTAATCAAAGCAGTGGCGTTGGCATGA 120

Query: 124 AGGCAGACAGAGGTCATGGAAGAGACCAGGCTCAGAAACAGCCCCACCATGCACAGCGGG 183
 |||
 Sbjct: 121 AGGCAGACAGAGGTCATGGAAGAGACCAGGCTCAGAAACAGCCCCACCATGCACAGCGGG 180

```

Query: 184 ATGTTTTCCCAAGGGCAACATGCAAAGCCAGGTATCCACATGGGTAGAGTAGAAAGT 243
      181 ATGTTTTCCCAAGGGCAACATGCAAAGCCAGGTATCCACATGGGTAGAGTAGAAAGT 240
5  Sbjct: 181 ATGTTTTCCCAAGGGCAACATGCAAAGCCAGGTATCCACATGGGTAGAGTAGAAAGT 240

Query: 244 CAGACCTTACATCTCACACAAATGAACTCAAATATACCAGAGAGCAAAGCTAAGAGC 303
      241 CAGACCTTACATCTCACACAAATGAACTCAAATATACCAGAGAGCAAAGCTAAGAGC 300
10 Sbjct: 241 CAGACCTTACATCTCACACAAATGAACTCAAATATACCAGAGAGCAAAGCTAAGAGC 300

Query: 304 TAAATCAAGTTTCTTAGGGCAAGCTGTAGTAGGCTCCCTTGGGTGGGTAAATGCTTTTG 363
      301 TAAATCAAGTTTCTTAGGGCAAGCTGTAGTAGGCTCCCTTGGGTGGGTAAATGCTTTTG 360
15 Sbjct: 301 TAAATCAAGTTTCTTAGGGCAAGCTGTAGTAGGCTCCCTTGGGTGGGTAAATGCTTTTG 360

Query: 364 TGGATGTGACTACCAAAAATTCAACCAGAGCCAACGACCCAACTATTAATGGGCAGTGGA 423
      361 TGGATGTGACTACCAAAAATTCAACCAGAGCCAACGACCCAACTATTAATGGGCAGTGGA 420
20 Sbjct: 361 TGGATGTGACTACCAAAAATTCAACCAGAGCCAACGACCCAACTATTAATGGGCAGTGGA 420

Query: 424 CCTAAAGAGATTCTTCAAACGATATATAAAGAAGGCCACCAAGCATATAAACATGTGA 483
      421 CCTAAAGAGATTCTTCAAACGATATATAAAGAAGGCCACCAAGCATATAAACATGTGA 480
25 Sbjct: 421 CCTAAAGAGATTCTTCAAACGATATATAAAGAAGGCCACCAAGCATATAAACATGTGA 480

Query: 484 CATCAGTAGTCAGAGAGATGGGAAGCAGAAGCACTAGCAGATCTTAACACCTACTAGAAC 543
      481 CATCAGTAGTCAGAGAGATGGGAAGCAGAAGCACTAGCAGATCTTAACACCTACTAGAAC 540
30 Sbjct: 481 CATCAGTAGTCAGAGAGATGGGAAGCAGAAGCACTAGCAGATCTTAACACCTACTAGAAC 540

Query: 544 AGCCACTAAAAAGAGTAAGACTCACAAGGACATGGGCACTTCTAATCTCTGTGCACTGC 603
      541 AGCCACTAAAAAGAGTAAGACTCACAAGGACATGGGCACTTCTAATCTCTGTGCACTGC 600
35 Sbjct: 541 AGCCACTAAAAAGAGTAAGACTCACAAGGACATGGGCACTTCTAATCTCTGTGCACTGC 600

Query: 604 TGCCAGGACATACAATAGTGTGGTCACTATGGAGACTACGGCAGTGCCTACTAATAACAG 663
      601 TGCCAGGACATACAATAGTGTGGTCACTATGGAGACTACGGCAGTGCCTACTAATAACAG 660
40 Sbjct: 601 TGCCAGGACATACAATAGTGTGGTCACTATGGAGACTACGGCAGTGCCTACTAATAACAG 660

Query: 664 C 664
      661 C 661

```

INJURYMARKER2

INJURYMARKER2 is a 893 nucleotide sequence encoding phosphatidylethanolamine N-methyltransferase [L14441]:

```

40 1 tccccgctga gttcatcacc agggacaggt gacctgagct gcccctggag cccagctccc
61 atttccttct ggttctggcc gatctcttcg ttatgagctg gctgctgggt tacgtggacc
121 ccacagagcc cagctttgtg gcggtctgtc tcaccattgt gttcaatcca ctcttctgga
181 atgtggttagc aaggtgggag cagagaactc gcaagctgag cagagccttc ggggtcccctt
241 acctagcctg ctattccctg ggcagcatca tcctgcttct gaacatcctc cgctcccact
45 301 gcttcacaca ggccatgatg agccagccca agatggaggg cctggatagc cacaccatct
361 acttcctggg ccttgcaactc ctgggctggg gactcgtggt tgtgctctcc agcttctatg
421 cactgggggtt cactggggacc tttctaggtg actacttttg gatcctcaag gagtccagag
481 tgaccacatt tccttcagc gtgctggaca accccatgta ctggggaagt acagccaact
541 acctaggctg ggcacttatg cacgccagcc ctacaggcct gctgttgacg gtgctggtgg
50 601 cactcgtcta cgtggttgct ctctgtttg aagagccctt cactgcggag atctaccggc
661 ggaaagccac caggttgcac aaaaggagct gacagggcca tgagggacct ttggaaagcc
721 ggattgcctc ccggtgacc caagcaaca cccttctcg ggagagcagc gctggccatt
781 gtacctgtgc cttggaaacc agtcatgggg gtgctcaggc attatgtcat gtgactgctg
841 agaccccat cccaccaat ccctgacaca ctaataaagg ctttgtgacc tcc

```

INJURYMARKER3

INJURYMARKER3 is a 1131 nucleotide hexokinase-encoding sequence [M86235]:

```

1 agcaggaatc ccctccgctt gcgggtagga agcttgggga gcagcctcat
60 ggaagagaag

```

```

61   cagatcctgt gcgtaggggt ggtggtgctg gacatcatca atgtggtgga
    caaataccca
121  gaggaagaca cggatcgagc gtgcctatcc cagagatggc agcgtggagg
    caacgcgtcc
5   181  aactcctgca ctgtgctttc cttgctcgga gcccgtgtg ccttcattgg
    ctgctggcc
    241  catggccatg ttgccgactt cctggtggcc gacttcaggc ggaggggtgt
    ggatgtgtct
    301  caagtggcct ggagagcca gggagatacc ccttgctcct gctgcatcgt
10   caacaactcc
    361  aatggctccc gtaccattat tctctacgac acgaacctgc cagatgtgtc
    tgctaaggac
    421  tttagaagg tcgatctgac ccggttcaag tggatccaca ttgagggccg
    gaatgcatcg
15   481  gaacaggtaa agatgctaca gcgtagataa cagtacaatg ccacgcagcc
    tctgcagcag
    541  aaggtccggg tgtccgtgga gatagagaag ccccgagagg aactcttcca
    gctgttcggc
    601  tatggagagg tgggttttgt cagcaaagat gtggccaagc acctgggggt
20   ccggtcagca
    661  ggggaggccc tgaagggtt gtacagtcgt gtgaagaaag gggctacgct
    catctgtgcc
    721  tgggctgagg agggagccga tgcctgggc cccgacggcc agctgctcca
    ctcagatgcc
25   781  tccccaccac cccgagtagt agacactctc ggggctggag acaccttcaa
    tgccctctgc
    841  atcttcagcc tctccaaggg aaacagcatg caggaggccc tgagattcgg
    gtgccagggtg
    901  gctggcaaga agtgtggctt gcaggggttt gatggcattg tgtgagagat
30   gagcgggtgg
    961  aggtagcagc tcgacacctc agaggctggc accactgcct gccattgcct
    tcttcatttc
    1021 atccagcctg gcgtctggct gccagttcc ctgggccagt gtaggctgtg
    gaacgggtct
35   1081 ttctgtctct tctctgcaga cacctggagc aaataaatct tcccctgagc c

```

INJURYMARKER4

INJURYMARKER4 is a 1994 nucleotide sequence encoding mitochondrial HMG-CoA Synthase [M33648]:

```

40   1   atctctccca ggggctgtgg actgctggct ttctgttgat accttagaga
    tgcagcggt
    61   tttggctcca gcaaggcggg tcttgcaagt gaagagagtc atgcaggaat
    cttcgtctc 121   acccgctcac ctgctccccg cagcccagca gaggtttct
    acaatccctc ctgctccct 181   ggccaaaact gatacatggc caaaagatgt
45   gggcatcctt gccctggagg tctactttcc 241   agcccaatat gtggacaaa
    ctgacctgga gaagttcaac aatgtggaag cagggaagta 301   cacagtgggc
    ttgggccaga ccggtatggg cttctgttcg gtccaggagg acatcaactc 361   cttgtgcctc
    acagtgggtc agaggctgat ggaacgcaca aagctgccat gggatgccgt 421   aggcgcctg
    gaagtgggca cggaaacat cattgacaag tccaaggctg tcaagacagt 481   gctcatggag
50   ctcttcagg attcaggcaa cactgacatc gagggcatag ataccaccaa 541   cgctgtctat
    ggtggcactg cctccctctt caacgctgcc aactggatgg agtccagcta 601   ctgggatggt
    cgctatgcc tgggtgtctg tgggtgatatc gcagtctacc caagtggtaa 661   cccccgccc
    acaggtggtg cgggggtgtg ggcaatgctg attgggcca aggccccgt 721   agtctggaa
    caagggctga ggggaacca catggagaac gcctatgact tctacaaacc 781   aaactggcc
55   tcagagtatc cactgggtgga tgggaagctg tctatccagt gctacctgcg 841   ggccttgga

```

c gatgctatg cagcttaccg caggaaaatc cagaatcagt ggaagcaagc 901 tggaaacaac
 cagcctttca ccctcgatga cgtgcaatat atgatcttcc acacaccctt 961 ttgcaagatg
 gtccagaaat ccctagctcg gctgatgttc aatgacttcc tgtcatctag 1021
 cagtgacaag cagaacaact tatacaaggg tctagaggcc ttcaagggtc
 5 taaagctgga 1081 agaaacctac accaacaagg atgttgacaa ggctctgctg
 aaggcctccc tggacatgtt 1141 caacaagaaa accaaggcct ccctttacct
 ctccacaaac aatgggaaca tgtacacctc 1201 gtccctctac ggggtgcctgg
 cctcacttct ctcccaccac tctgcccagg aattggccgg 1261 ctccaggatt
 ggagccttct cctacggctc aggttagca gcaagtttct tctcatttct 1321
 10 agtgtccaag gacgttccc caggttccc tctggagaag ctggtgtcta
 gtgtgtcaga 1381 tctgcccaga cgtctagact cccggagacg catgtccct
 gaggaattca cagaaataat 1441 gaatcagaga gagcaatttt accacaaggt
 gaacttctct cccctggtg acacaagcaa 1501 cctcttccca ggcacttggg
 accttgaacg agtggatgag atgcaccgca gaaaatatgc 1561 ccggcgctccc
 15 gtctaaggag accaatccat acaaccattc cccggggaaa gaatgtgagc 1621
 agagccgtta cccaaacggc ttccacttaa aattccacc acagcagtga
 acggtgaata 1681 gacacagcga ccccatagga tctgctccgc ggtgaagggc
 ctccctctgt ggatcctggg 1741 tgaccctccc tgaagcagtg agcaccacag
 gttctgctgt ggaccagagc cccctgtgg 1801 agaggagaa agaaagggga
 20 gccgctgacc tgcagggata cagaccttcc ccacagcctg 1861 gcagccgccc
 gtttgttga gcttattatc agactgtggg ctatcatagt tcatgctcgt 1921
 ttcttaaagt ttcccagaaa ttctaaaaat tttgtatcta aacttttaat
 atggcgatta 1981 aaaggagaga agga

25 INJURYMARKER5

INJURYMARKER5 is a 1850 nucleotide sequence encoding cathepsin C [D90404],
 having the following nucleic acid sequence:

1 gaattccggt tctagttgtt gttttctctg ccactctgctc tccgggcgcc gtcaaccatg
 61 ggtccgtgga cccactcctt gcgcgcgcgc ctgctgctgg tgcttttggg agtctgcacc
 30 121 gtgagctccg acactcctgc caactgcact taccctgacc tgctgggtac ctgggtttc
 181 caggtggggc ctagacatcc ccgaagtcac attaaactgct cggtaatgga accaacagaa
 241 gaaaaggtag tgataacct gaagaagttg gatactgcct atgatgaagt gggcaattct
 301 gggattttca cctcattta caaccaaggc ttgagattg tgttgatga
 ctacaagtgg 361 ttgctgtttt tcaagtatga agtcaaaggc agcagagcca
 35 tcagttactg ccattgagacc 421 atgacagggt ggggccatga tgcctggggc
 cggaactggg cttgctttgt tggcaagaag 481 atggcaaate actctgagaa
 ggtttatgtg aatgtggcac accctggagg tctccaggaa 541 aaatattctg
 aaaggctcta cagtcacaac cacaactttg tgaaggccat caattctgtt 601 cagaagtctt
 ggactgcaac cacctatgaa gaatatgaga aactgagcat acgagatttg 661 ataaggagaa
 40 gtggccacag cggaaggatc ctaaggccca aacctgcccc gataactgat 721 gaaatacagc
 aacaaatttt aagtttgcca gaatcttggg actggagaaa cgtccgtggc 781 atcaattttg
 ttagccctgt tcgaaaccaa gaatcttgtg gaagctgcta ctcatattgcc 841 tctctgggta
 tgctagaagc aagaattcgt atattaacca acaattctca gacccaate 901 ctgagtcctc
 aggaggttgt atctttagc ccgtatgcc aaggttgtga tgggtgattc 961 ccatacctca
 45 ttgcaggaaa gtatgcccga gattttgggg tgggtggaaga aaactgcttt 1021
 ccctacacag ccacagatgc tccatgcaaa ccaaaggaaa actgcctccg
 ttactattct 1081 tctgagtact actatgtggg tggtttctat ggtggctgca
 atgaagccct gatgaagctt 1141 gagctgggtca aacacggacc catggcagtt
 gcctttgaag tccacgatga ctctctgcac 1201 taccacagtg ggatctacca
 50 ccacactgga ctgagcgacc ctttcaacc ctttgagctg 1261 accaatcatg
 ctgttctgct tgtgggctat ggaaaagatc cagtcactgg gtagactac 1321
 tggattgtca agaacagctg gggctctcaa tggggtgaga gtggctactt
 ccggatccgc 1381 agagggaactg atgaatgtgc aattgagagt atagccatgg
 cagccatacc gattcctaaa 1441 ttgtaggacc tagctcccag tgtccatac
 55 agctttttat tattcacagg gtgatttagt 1501 cacaggctgg agacttttac
 aaagcaatat cagaagctta ccactaggtta cccttaaga 1561 attttgcctt

taagtttaaa acaatccttg atttttttct ttttaatatcc tccctatcaa 1621
 tcaccgaact acttttcttt ttaaagtact tggtaagta atacttttct
 gaggattggt 1681 tagatattgt caaatatttt tgctgggtcac ctaaaatgca
 gccagatggt tcattgttaa 1741 aaatctatat aaaagtgcaa gctgcctttt
 5 ttaaattaca taaatcccat gaatacatgg 1801 ccaaaatagt tattttttaa
 agactttaaa ataaatgatt aatcgatgct

INJURYMARKER6

INJURYMARKER6 is a 993 nucleotide sequence encoding hydroxysteroid

10 sulfotransferase [D14989]:

1	ggcaagggct	ggaatactaa	aagttattca	tgatgtcaga	ctatacttgg	tttgaaggaa
61	taccttttcc	tgcccttttg	ttttccaaag	aaattctgga	aaatagttgt	aagaagtttg
121	tggtaaaaga	agacgacttg	atcatattga	cttaccctaa	gtcaggaacg	aactggctga
181	tcgagattgt	ctgcttgatt	cagaccaagg	gagatcccaa	gtggatccaa	tctatgccc
15 241	tctgggatcg	ctcaccctgg	atagagactg	gttcaggata	tgataaatta	accaaattgg
301	aaggaccacg	actcatgacc	tcccatcttc	ccatgcatct	tttctccaag	tctctcttca
361	gttccaaggc	caaggtgata	tatctcatca	gaaatcccag	agatgttctt	gtttctgctt
421	atthttttctg	gagtaagatc	gccttgagga	agaaaccaga	ctcgctggga	acttacgttg
481	aatggttcct	caaaggaaat	gttgcatatg	gatcatgggt	tgagcacatc	cgtggctggc
20 541	tgtctatgag	agaatgggac	aacttcttgg	tactgtacta	tgaagacatg	aaaaaggata
601	caatgggacg	cataaagaag	atatgtgact	tcctggggaa	aaaattagag	ccagatgagc
661	tgaatttggt	cctcaagtat	agttccttcc	aagtcgtgaa	agaaaacaac	atgtccaatt
721	atagcctcat	ggagaaggaa	ctgattctta	ctggttttac	tttcatgaga	aaaggcacia
781	ctaagtactg	gaagaatcac	ttcacagtag	cccaagctga	agcctttgat	aaagtgttcc
25 841	aggagaaaat	ggccgggttc	cctccaggga	tggtcccatg	ggaataaatt	ttcaaaagtt
901	ttaaattattt	tatgaacact	gatgtttatg	tttatgttgt	tctatgatgt	ctgaataact
961	gaatgtgatc	attgaataaa	tcctgttgtg	gat		

INJURYMARKER7

30 INJURYMARKER7 is a 5001 nucleotide sequence encoding insulin-like growth factor
 binding protein [L22979]:

1	cacaaaccca	gcgagcattg	aacactgcac	acggccatct	gcccagagag	ctgtgaccac
61	cacttccgct	actatctact	cagaaagtcg	tgactactga	gccactgctg	cctgcccaga
121	ttctcatcca	ccgcctgctg	cgtctgggtg	cgatgccgga	gttcctaact	gttgtttctt
35 181	ggccgttccct	gatcctcctg	tccttccagg	ttcgcgtagt	cgctggagcc	ccccagccat
241	ggcaactgtgc	tccttgcact	gctgagaggc	tgaggtctctg	tccaccctg	cctgcttcgt
301	gccccgagat	ttctcggcct	gcgggctgtg	gctgctgccc	gacatgtgcc	ttgccactgg
361	gtgctgcctg	tggtgtggcc	actgcggcct	gcgctcagg	actcagctgc	cgtgcgctgc
421	caggggagcc	tcgacctctg	catgccctca	cccgtggcca	gggagcctgt	gtactagaac
40 481	ctgccgcacc	cgccacgagc	agcttgctcg	gttctcagca	tgaaggtagt	acagccctct
541	ctgcctcttg	atctcttggc	taggacacac	gtgctttcta	ggcacgtcag	aggcctatcc
601	ggaacctata	gcagatagga	caaaggtctt	ccatgcccac	tttgagcttt	cagcctcaaa
661	taaggccctc	agttaggtcg	tggcggtctg	ggaacaccca	gaggtgtcaa	tccagtagca
721	gagtggaaga	gttgggaaga	atgttccaag	ctcccagtg	agagtggaga	gttgggaaga
45 781	atgtctcacag	actaggtagt	actgatcctg	cttggtcttt	cagtggggag	ggagctatgg
841	ggctgccagg	tgggtggggg	gctggcccaa	acacctcttt	ctgtgggtcc	tgaccttggc
901	agttccaatg	gctaaaaggt	ccaggaaggt	ttaggatggg	agccctcctg	ctgccccag
961	gaggtttgca	atgtcctttg	tagcatatat	cctgccacac	agtatgtgct	tcccagatgt

	1021	ttacagaaca	taatgtgaaa	atttaggccc	aaaccttcac	ttccattcat	tgctatagac
	1081	aaacagtgtt	tgaagtgtat	gttgccctgct	aggagtctga	caatcaggcg	ctttcctgaa
	1141	tttaagcact	ggtttggttg	taataggaag	cttgggaaat	gcctcttcct	ctgctccagc
	1201	ccctatctcc	cctgtctggg	ctgcatgcac	ttcctgtgtg	ggtaaggagc	ctcatggttc
5	1261	catattctga	cggaagccg	gactgcaggc	atctgacctt	tttgactaaa	tggaagaact
	1321	atcccaacgg	tccttagaaa	cgggcttccc	caggagcgat	gtctgataat	gtcctcctct
	1381	gtgaggggct	gcctaagagg	tgctcgggtt	caagaaagca	gggctcccag	aaaagaagag
	1441	gatgggtggt	tgaggtgggg	aaggtacac	tctacacctt	gcttctcaac	tatcccccta
	1501	ctggggctct	acgagattct	ttttgtggtg	tggagaggag	agctgagtgg	tcaagttctca
10	1561	ccactaacgg	gttcaagcct	tggcctcagt	ccttggcttc	ttcaggatta	catccctagac
	1621	ccaactctct	ctgccatggg	gactcccttg	cctaacccca	aaacatacca	tttccccaga
	1681	aaggaattag	tattgctaata	tggtgataat	tggtcccaaa	tagccactg	tgaaaacaa
	1741	agcctgattt	cacctgactg	ttacagattg	gtcttaaggc	ggtagacgtg	agtgcacatag
	1801	gagtgcacac	tcagggtctca	tcgtctgtgt	ctgtgggggt	cgttttcaga	ggcaaagggt
15	1861	gctgtggcct	ctgaggatga	gcttgccgag	agcccagaga	tgacagagga	acagctgctg
	1921	gatagcttcc	acctcatggc	ccatcccgt	gaggaccagc	ccatcctgtg	gaatgccatt
	1981	agcacctaca	gcagcatgcg	ggcccgagg	atcactgacc	tcaagaaatg	gaaggtgaga
	2041	ccctgcactc	agaccttcag	gttttagctat	ctacgtgaag	aggtttgtct	agacgatttc
	2101	ttaaagggca	ctgagcatgg	ggctgagaac	ggggatataa	ctacccccat	ccctgatgta
20	2161	tttctgcctc	cttaaaaaata	tggcaagtat	ctcagagcat	aaggtagggc	atttttcagt
	2221	ctagggtttct	ctgtcaccga	gtacgcacgt	tcagtgtatt	ttagccacca	accagctcca
	2281	cggttttgcc	agccttttagc	tatgcacttt	agctatgcag	taaactttct	tagctttact
	2341	ggctgttttt	caacttgacc	acttggggag	gacagagaac	caaagggtga	gagaaagtac
	2401	ggcagaagca	ttgaagaagt	acaccttaag	aaatgaaagg	ataaacattg	ttagggtcac
25	2461	tttagaattt	catatggaaa	ttgtccaaat	cagtgccttg	ttccgtaatc	aatttgacat
	2521	acaccaaagt	caaggatggc	tgtttgaaaa	atctaggcat	ttatgatgct	aaattccaca
	2581	cacagagact	gagcctgtct	tttttattag	agttcagggt	ctcaagttat	tcagagatag
	2641	ccagggtcag	gaagcattta	taccattggc	caggctctta	ccacaatgtc	gttaaggggg
	2701	tctccagaaa	atgccactga	gggaggatga	gagtgggtgc	cctgtccttt	atctacatag
30	2761	cccaagccag	agaccaacct	gtcctgtctca	cagatgggga	aacatctcag	ccgttgtcta
	2821	aattgataat	ttttgtctct	gttactcatg	ctaataataa	attatccttt	taggagccct
	2881	gccaacggga	actctataaa	gtgttagaga	gattagctgc	cgctcaacag	aaagcaggag
	2941	atgagatcta	caaattttat	ctgccaaact	gcaacaagaa	tggtttttat	cacagcaaac
	3001	aggtaggtgg	ctttgtctcat	ccagatcctt	gtaaaacttc	atgatttttt	tttttaaagt
35	3061	caaatgattc	acaggcccaa	tacacatcat	gggtagcttt	cttaggtgag	atccagccct
	3121	gcagtagtgt	ggagaagcta	gtcctgagaa	agagatagt	tgatggatga	ggaacacttc
	3181	agccagaagg	gaggactaag	cattagtgtg	atgagtggag	agcacttcag	ttaacaggga
	3241	ggactaagca	ttagtgtgat	gagtggagac	cacttcaagc	cagagggagg	actaacattg
	3301	gcagtatgat	gagtggagg	cacttcagcc	agtagggagg	actaaccatt	agtctcatca
40	3361	ctaggagcac	ctcagccaag	tagggaggac	taaccattag	tctcacactc	acccaacatc
	3421	ttcagtccag	actaagcatt	agtgtgatga	gtgaggagca	cttcagtcag	tagggaggac
	3481	tacattagt	tgatgagtga	ggagcacttc	agccagtagg	gaggactaac	cgttcactca
	3541	gattagcaga	gatggatgtt	ccatatactg	atgtccagg	ttcagttcct	cacaactaga
	3601	ggaaagggac	acagtcagtg	taggagacag	atgtctcgcg	ttctctcttc	ccacaaataa
45	3661	aaacaaactc	tgtagtaaga	cacaccaatt	gtgctttgcc	tagcaataaa	tgagattgaa
	3721	gaagtccagg	cttaattttcg	acgcaacttt	agaactcagg	gaagtgcagg	ttctggaatt
	3781	tcattgagga	aaaacttgag	gtctaggtct	agccgtgtgg	tagagatggg	gagacctatc
	3841	gttgagctcc	tttggcagag	ggccatggag	caggtaaccg	tcaaaacaat	ataccactga
	3901	gtaaacagat	gagattgtta	tcagggtgtg	cataaagcca	acctctccgt	tttgtgatga
50	3961	caaccagaag	ggcattggtc	tgccgagcct	tagccagcag	gtagctgtgc	agtgtttggc
	4021	ctcactgagg	gacaggggtg	ccagagctct	tacctcctgc	tgctcttgac	ctcgttctg
	4081	tctttgcagt	gcgagacatc	tctggatgga	gaagctgggc	tctgctgggt	tgtctaccca
	4141	tgagtgggga	agaagatccc	tggtctctct	gagaccagag	gggaccccaa	ctgccaccag
	4201	tatttttaagt	tgcaaaactg	aaagttgttt	cctccctcct	tcttcacaca	aaatatttta
55	4261	gtatatagt	tatttatact	cgggagcaca	ccattttata	tatgtgtata	tgtatatatc
	4321	caggaaactag	tttttatact	ccacatgctg	cttgatgtac	aagtgggttt	gtatttattc
	4381	actctaagtt	tatttttttc	taccctgtcc	ttgtgctgta	tttaatttata	taactgaagc
	4441	ttttctcatc	tccatacatg	taaaactac	catctcagct	cttccagagt	tctgctttga
	4501	aagggtcagc	cggtacgtgc	ctagaacag	cacaagtcag	tctgaggtag	gggcttttca
60	4561	gtgggttcag	ggaggaaggt	tagccctggc	tcggggagac	ttcctcatcg	aatccacag
	4621	gtctgtgtct	gatgcctatt	ggctgggaag	gttccgatgt	tggttgtgta	atcaaagcta
	4681	aacgtggaaa	gctgcgtccc	atgcactgtt	aaacacacgt	ctggaataaa	acattctacc
	4741	tggaaacact	gctgtctctg	tggaattcca	gctctgtgct	cattccctca	gtccgttcgg

4801 ctttcccgcgct cgccctgattc ctgggtctgt gctttgggga tagatgttgc aatacaggggt
 4861 gcttggttgt ttacagaaca ccctggacaa acactctgtg actttatggt cccattttca
 4921 agcagcatca ggcctctgtc tgggccagac tacagagccc ctccctcctg gtccatctcc
 4981 ctttcttccc agggccctca g

5

INJURYMARKER8

INJURYMARKER8 is 579 bp rat expressed sequence tag (EST) [AA851963]. The nucleic acid was initially identified in a cloned fragment having the following sequence:

1 TGTACATAATTTATTAATAATGTCTCTGACACAAATAATGACTCCACTGCATACATAGTTGGTGTTCAAAAATTTCCCCA
 81 ATGTTTGTCTGGACACAATTGTTATTAGCCAACTCGGTGAATTCAGACATTGTTCCACACAATGAACAATCGCACACA
 161 TGAGAACTGCACCTAGAATGTCCATCCTAGAATCTCCATCCATCCAGTCAAAGTGCTGAGCTCACTGACTGAAGGAAACA
 241 TGACCTGTGTTCTAGA (SEQ ID NO: 13)

10 The cloned sequence was assembled into a contig resulting in the following consensus sequence:

1 TGTACATAATTTATTAATAATGTCTCTGACACAAATAATGACTCCACTGCATACATAGTTGGTGTTCAAAAATTTCCCCA
 81 ATGTTTGTCTGGACACAATTGTTATAAGCCAACTCGGTGAATTCAGACATTGTTCCACACAATGAACAATCGCACACA
 161 TGAGAACTGCACCTAGAATGTCCATCCTAGAATCTCCATCCATCCAGTCAAAGTGCTGAGCTCACTGACTGAAGGAAACA
 241 TGACCTGTGTTCTAGAACGTAGCTGGCTATGAAGTTACTCATGTGTAAATTCCTTAAAAAGATTAAATTGTTTGGCCCA
 321 TTTCTATATTTTCAATAAATACTATAATTACAACTTTCTAAAAATAATTTTACAACCATGTAATTATGACTAACCATAT
 401 CATCTAAAAAGTAAGTGAAGTCATTGTCTAGAGATTGTCTGAGATTATTCTGCTGAGAAGCTTACTTCAAACCTCTTATC
 481 ACTACTTCTACTTCCAGTGTCTTGAATTAAGAACAGAAATTGTAACATATGCTATTCTACATCAGATTGACACAACCTA
 561 CTTCTAAGTACACTATTGC (SEQ ID NO: 14)

Blast-N Results:

>gb:GENBANK-ID:AA851963|acc:AA851963 EST194732 Normalized rat spleen, Bento Soares
 Rattus sp. cDNA clone RSPAO86 3' end, mRNA sequence - Rattus sp., 538 bp (RNA).

Top Previous Match Next Match
 Length = 538

15 Plus Strand HSPs:

Score = 2681 (402.3 bits), Expect = 8.1e-115, P = 8.1e-115
 Identities = 537/538 (99%), Positives = 537/538 (99%), Strand = Plus / Plus

20 Query: 42 CTCCACTGCATACATAGTTGGTGTTCAAAAATTTCCCAATGTTTGTCTGGACACAATT 101
 |||
 Sbjct: 1 CTCCACTGCATACATAGTTGGTGTTCAAAAATTTCCCAATGTTTGTCTGGACACAATT 60

25 Query: 102 GTTATAAGCCAACTCGGTGAATTCAGACATTGTTCCACACAATGAACAATCGCACACAT 161
 |||
 Sbjct: 61 GTTATTAGCCAACTCGGTGAATTCAGACATTGTTCCACACAATGAACAATCGCACACAT 120

Query: 162 GAGAACTGCACCTAGAATGTCCATCCTAGAATCTCCATCCATCCAGTCAAAGTGCTGAGC 221

```

Sbjct: 121 |||||GAGAACTGCACCTAGAAATGTCCATCCTAGAAATCTCCATCCATCCAGTCAAAGTGCTGAGC 180
5 Query: 222 TCACTGACTGAAGGAAACATGACCTGTGTTCTAGAACGTAGCTGGCTATGAAGTTTACTC 281
Sbjct: 181 TCACTGACTGAAGGAAACATGACCTGTGTTCTAGAACGTAGCTGGCTATGAAGTTTACTC 240
Query: 282 ATGTGTAAATTCCCTTAAAAAGATTAAATTGTTTGGCCCATTTCTATATTTTATAAAATAA 341
10 Sbjct: 241 ATGTGTAAATTCCCTTAAAAAGATTAAATTGTTTGGCCCATTTCTATATTTTATAAAATAA 300
Query: 342 CTATAATTACAAACTTTCTAAAAATAATTTTACAACCATGTAATTATGACTAACCATATC 401
Sbjct: 301 CTATAATTACAAACTTTCTAAAAATAATTTTACAACCATGTAATTATGACTAACCATATC 360
15 Query: 402 ATCTAAAAAGTAAGTGAAGTCATTGTCTCTAGAGATTGTCTGAGATTATTCTGCTGAGAAG 461
Sbjct: 361 ATCTAAAAAGTAAGTGAAGTCATTGTCTCTAGAGATTGTCTGAGATTATTCTGCTGAGAAG 420
20 Query: 462 CTTACTTCAAACCTCTTATCACTACTTCTACTTCCAGTGTCTTGAATTAAGAACAGAAA 521
Sbjct: 421 CTTACTTCAAACCTCTTATCACTACTTCTACTTCCAGTGTCTTGAATTAAGAACAGAAA 480
Query: 522 TTGTAACATATGCTATTCTACATCAGATTGACACAACCTACTTCTAAGTACACTATTGC 579
25 Sbjct: 481 TTGTAACATATGCTATTCTACATCAGATTGACACAACCTACTTCTAAGTACACTATTGC 538

```

INJURYMARKER9

INJURYMARKER9 is a 2495 nucleotide catalase-encoding sequence[M11670],

30 having the following nucleic acid sequence:

```

1 attgcctacc ccgggtggag accgtgctcg tccggccctc ttgcctcacg ttctgcagct
61 ctgcagctcc gcaatccctac accatggcgg acagccggga cccagccagc gaccagatga
121 agcagtgga ggaacagcgg gccctcaga aacccgatgt cctgaccacc ggaggcggga
181 acccaatagg agataaactt aatatcatga ctgcggggcc ccgagggccc ctctcgttc
35 241 aagatgtggt ttaccgcgac gagatggcac actttgacag agagcggatt cctgagagag
301 tggtagatgc aaagggagca ggtgcttttg gatactttga ggtcaccac gatattacca
361 gatactccaa ggcaaagggtg tttgagcata ttgggaagag gactcctatt gccgtccgat
421 tctccacagt cgctggagag tcaggctcag ctgacacagt tcgtgacct cgtgggtttg
481 cagtgaatct ctacactgaa gatggttaact gggacctcgt gggaaacaac acccctattt
40 541 tcttcatcag gtagccatg ttgtttccat cctttatcca tagccagaag agaaaccac
601 aaactcacct gaaggacct gacatggtct gggacttctg gactctttgt ccagagtctc
661 tccatcaggt tactttcttg ttcagcgacc gagggattcc agatggacat cggcacatga
721 atggctatgg ctacacacac ttcaagctgg ttaatgagaa tggagaggca gtgtactgca
781 agttccatta caagactgac cagggcatca aaaacttgcc tgttgaagag gcaggaagac
45 841 ttgcacagga agaccggat tatggcctcc gagatctttt caatgccatc gccagtggca
901 attaccatc ctggactttt tacatccagg tcatgacttt caaggaggca gaaaccttcc
961 catttaatcc atttgacctg accaagggtt ggctcacaac ggactacctt cttataccag
1021 ttggcaaact ggtcttaaac agaaatcctg ctaattatct tgcgtgaagt gaacagatgg
1081 cttttgaccc aagcaacatg ccccttgga ttgagcccag cccggacaag atgtccagg
50 1141 gccgcctttt tgcttaccca gacactcacc gccaccgctt gggaccaaac tatctgcaga
1201 tacctgtgaa ctgtccctac cgtgctcgcg tggccaacta ccagcgcgat ggccccatgt
1261 gcatgcatga caaccagggt ggtgctccca actactacc caacagcttc agcgaccag
1321 agcagcagg ctcggccctg gagcaccata gccagtgtct tcagatgtg aagcgcttca
1381 acagtgttaa tgaagacaac gtcactcagg tgcggacatt ctatacgaag gtgttgaatg
55 1441 aggaggagag gaaacgcctg tgtgagaaca ttgccaacca cctgaaagat gctcagcttt
1501 tcattcagag gaaagcggtc aagaatttca ctgacgtcca cctgactac ggggcccgag
1561 tccaggctct tctggaccag tacaactccc agaagcctaa gaatgcaatt cacacctacg
1621 tacaggccgg ctctcacata gctgccaaag gaaaagctaa cctgtaaagc acgggtgctc
1681 agcctcctca gcctgactg aggagatccc tcataagca gggcacaagc ctcaccagta
60 1741 atcatcgctg gatggagtct cccctgctga agcgagact cactgtgacg tctttaaac
1801 gataatccaa gcttctagag tgaatgatag ccatgctttt gatgacattt cccgaggggg
1861 aaattaaaga ttagggctta gcaatcactt aacagaaaca tggatctgct taggacttct
1921 gtttggatta ttcatttaaa atgattacaa gaaagggttt ctagccagaa acatgatttg

```

```

1981  attagatatg  atatatgata  aaatcttggg  gattttacta  tagtcttatg  ttacctcaca
2041  gcctgggtata  tataacaacac  acacacacac  acacacacac  acacacacaaa  acacacatac
2101  actatacaca  cacacacaca  cacacacata  aacacacata  cacaacacac  acatacacta
2161  cacacacaga  acacacaaca  caaacatata  cacataggca  cacacacaca  cacacacaca
5    2221  cacacacaca  cacacacaca  cacacatgaa  tgaagggatt  ataaagatgg  cccaccacaga
2281  attttttttt  atttttctaa  ggtccttata  agaaaaacca  tacttggatc  atgtcttcca
2341  aaaataactt  tagcactggt  gaaacttaat  gtttattcct  gtgtagttga  ttggattcct
2401  tttccccttg  aaattatggt  tatgctgata  cacagtgatt  tcacataggg  tgatttqtat
2461  ttgcttacat  ttttacaata  aaatgatcct  catgg

```

10

INJURYMARKER10

INJURYMARKER10 is a 1884 nucleotide betaine homocysteine methyl transferase-encoding sequence [AF038870]:

```

1    caagcctttg  ctggagaccg  ctctgttcca  gtccgcagct  ggcttcagcg  ccactcagga
15  61    caccggaaag  atggcaccga  ttgccggcaa  gaaggccaag  aggggaatct  tagaacgctt
121  aaatgctggc  gaagtcgtga  tcggagatgg  gggatttgct  ttgacttg  aaaagagggg
181  ctacgtaaa  gctggacct  ggacccaga  ggctgcggtg  gagaccccg  aggcagttcg
241  gcagcttcat  cgggagttcc  tcagagctgg  atcgaacgtc  atgcagacct  tcactttcta
301  tgcaagtga  gacaagctgg  aaaaccgagg  gaactacgtg  gcagagaaga  tatctgggca
20  361  gaaggtcaat  gaagctgctt  gtgacattgc  acggcaagtt  gctgacgaag  gggatgcatt
421  ggttgcagga  ggtgtgagtc  agacaccttc  ctacctcagc  tgcaagagtg  agacggaagt
481  taaaaagata  tttcaccaac  agcttgaggt  cttcatgaag  aagaatgtgg  acttctcat
541  tgcagagtat  tttgaacatg  ttgaagaagc  cgtgtgggca  gtcgaggcct  taaaaacatc
601  cgggaagcct  atagcggcta  ccattgtgat  cggacctgaa  ggagatctac  atggcggtgc
25  661  tcttgagag  tgcgcagtg  gtttggtaaa  agcaggtg  gccattgtcg  gtgtgaactg
721  ccacttcgac  cccagcacca  gcttcgagac  aataaagctc  atgaaggagg  gtctggaagc
781  agctcggctg  aaggcttact  tgatgagcca  cgccctggcc  taccacaccc  ctgactgtgg
841  caaacagga  tttattgatc  tcccagaatt  ccccttggga  ttggaaccca  gagttgccac
901  cagatgggat  attcaaaaat  acgccagaga  ggcctacaac  ctgggggtca  ggtacattgg
30  961  cggctgctgc  ggatttgagc  cctaccacat  cagggccatt  gcagaggagc  tcgcccaga
1021  aaggggattt  ttaccaccag  cttcagaaaa  acatggcagc  tggggaagtg  gtttgacat
1081  gcacacacaaa  ccctggatca  gggcaagggc  caggaaagaa  tactggcaga  atcttcgaat
1141  agcttcgggc  agaccgtaca  atccttcgat  gtccaagccg  gatgcttggg  gagtacgaa
1201  aggggcagca  gagctgatgc  agcagaagga  agccaccact  gagcagcagc  tgagagcgct
35  1261  cttcgaaaaa  caaaaattca  aatccgcaca  gttagccacag  gccagcggtt  cggggcgaa
1321  tctccagggt  ccgggccaca  gtgtgcacc  ggaaggagaa  ggcattctta  aaccagcggt
1381  tgtgttgatg  ccggttaca  cctgtgattg  gtgctagtta  gacaaaatgg  agtcacagat
1441  agcatttcac  agttacaaaa  ctacgcttta  gaattttacc  tagaaggaa  aaaggagaag
1501  tccacagtaa  atcctgaaca  catttcctac  gtgcctgtcg  cattacaggc  gcacaggagt
40  1561  cactgcagcg  aagagaaagt  caccgcagct  caatctcatt  tcagataggg  ggataggaca
1621  ccacctccac  gagtgacata  gaaccattca  gggaccgtat  cataagtgc  acagcaacca
1681  tctatatcta  agatgcttcc  caagtggatt  ccaagatcct  ttgagcagga  cccttaggca
1741  gaaacaacac  acaccagccc  tgtaaaactt  aacagataac  tgatccattc  tgtaattctg
1801  taatctctgt  tctgactgct  tccattccat  ttcattaata  aaaacatgcc  ggttgaaaaa
45  1861  cttcaaaaaa  aaaaaaaaaa  aaaa

```

Principle components analysis was used to generate three eigenvectors used to transform the original expression level data matrix, as shown in Table 6 below. Eigenvector 1 values represent NSAIDs associated with hepatotoxicity involving hepatocellular damage, Eigenvector 2 values represent NSAIDs associated with hepatotoxicity involving cholestasis,

and Eigenvector 3 values represent NSAIDs associated with hepatotoxicity involving elevated transaminase level.

Table 6: Transform Eigenvectors for Hepatotoxicity by Injury Type

Gene	Eigenvector 1	Eigenvector 2	Eigenvector3
INJURYMARKER1	58.7	0.325	-15.2
INJURYMARKER2	20.5	-3.23	3.01
INJURYMARKER3	-16.9	-6.52	-2.09
INJURYMARKER4	-10.3	0.351	-1.45
INJURYMARKER5	-7.59	-0.152	-0.310
INJURYMARKER6	11.4	-2.69	2.49
INJURYMARKER7	-16.0	-1.57	8.71
INJURYMARKER8	-11.6	1.13	5.36
INJURYMARKER9	-11.0	-0.351	0.078
INJURYMARKER10	7.55	0.618	4.65
% of variation explained	99.0	0.7	0.3

These eigenvectors may be used to transform the expression levels of INJURYMARKERS 1-10 ("INJURYMARKERS") in response to a given drug, in order to predict that drug's hepatotoxicity injury type. For example, expression levels of INJURYMARKERS correlating with Eigenvector 1 indicates that the test drug has a risk of hepatotoxicity involving hepatocellular damage. Alternatively, a drug's INJURYMARKERS expression profile can be generated simultaneously with the above-described training set (or an equivalent set) run in parallel with the test drug, and expression levels associated with the test drug directly compared to those of the training set.

GENERAL METHODS

The RISKMARKER (*i.e.* RISKMARKERS 1-8) and INJURYMARKER (*i.e.* INJURYMARKERS 1-10) nucleic acids and encoded polypeptides can be identified using the information provided above. In some embodiments, the RISKMARKER or INJURYMARKER nucleic acids and polypeptide correspond to nucleic acids or polypeptides which include the various sequences (referenced by SEQ ID NOs) disclosed for each RISKMARKER or INJURYMARKER polypeptide.

In its various aspects and embodiments, the invention includes providing a test cell population which includes at least one cell that is capable of expressing one or more of the sequences RISKMARKER 1-8 or INJURYMARKER 1-10. By "capable of expressing" is meant that the gene is present in an intact form in the cell and can be expressed. Expression of

one, some, or all of the RISKMARKER or INJURYMARKER sequences is then detected, if present, and, preferably, measured. Using sequence information provided by the database entries for the known sequences, or the sequence information for the newly described sequences, expression of the RISKMARKER or INJURYMARKER sequences can be
5 detected (if present) and measured using techniques well known to one of ordinary skill in the art. For example, sequences within the sequence database entries corresponding to RISKMARKER or INJURYMARKER sequences, or within the sequences disclosed herein, can be used to construct probes for detecting RISKMARKER or INJURYMARKER RNA sequences in, *e.g.*, northern blot hybridization analyses or methods which specifically, and,
10 preferably, quantitatively amplify specific nucleic acid sequences. As another example, the sequences can be used to construct primers for specifically amplifying the RISKMARKER or INJURYMARKER sequences in, *e.g.*, amplification-based detection methods such as reverse-transcription based polymerase chain reaction. When alterations in gene expression are associated with gene amplification or deletion, sequence comparisons in test and reference
15 populations can be made by comparing relative amounts of the examined DNA sequences in the test and reference cell populations.

Expression can be also measured at the protein level, *i.e.*, by measuring the levels of polypeptides encoded by the gene products described herein. Such methods are well known in the art and include, *e.g.*, immunoassays based on antibodies to proteins encoded by the genes.

20 Expression level of one or more of the RISKMARKER or INJURYMARKER sequences in the test cell population, *e.g.* rat hepatocytes, is then compared to expression levels of the sequences in one or more cells from a reference cell population. Expression of sequences in test and control populations of cells can be compared using any art-recognized method for comparing expression of nucleic acid sequences. For example, expression can be
25 compared using GENE CALLING® methods as described in US Patent No. 5,871,697 and in Shimkets et al., Nat. Biotechnol. 17:798-803.

In various embodiments, the expression of one or more sequences which are markers of hepatotoxicity risk, *i.e.* RISKMARKERS 1-8, is compared. In other embodiments, the expression of one or more sequences which are markers of hepatotoxicity injury type, *i.e.*
30 INJURYMARKERS, is compared. In further embodiments, expression of one or more RISKMARKERS and INJURYMARKERS may be compared to predict both hepatotoxicity risk and type of hepatotoxicity injury.

In various embodiments, the expression of 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, or all of the sequences represented by RISKMARKER 1-8 and INJURYMARKER 1-10 are measured. If desired, expression of these sequences can be measured along with other sequences whose expression is known to be altered according to one of the herein described parameters or conditions.

The reference cell population includes one or more cells for which the compared parameter is known. The compared parameter can be, *e.g.*, hepatotoxic agent expression status. By "hepatotoxic agent expression status" is meant that it is known whether the reference cell has had contact with a hepatotoxic agent. An example of a hepatotoxic agent is, *e.g.*, a thiazolidinedione such as troglitazone. Whether or not comparison of the gene expression profile in the test cell population to the reference cell population reveals the presence, or degree, of the measured parameter depends on the composition of the reference cell population. For example, if the reference cell population is composed of cells that have not been treated with a known hepatotoxic agent, a similar gene expression level in the test cell population and a reference cell population indicates the test agent is not a hepatotoxic agent. Conversely, if the reference cell population is made up of cells that have been treated with a hepatotoxic agent, a similar gene expression profile between the test cell population and the reference cell population indicates the test agent is a hepatotoxic agent.

In various embodiments, a RISKMARKER or INJURYMARKER sequence in a test cell population is considered comparable in expression level to the expression level of the RISKMARKER or INJURYMARKER sequence if its expression level varies within a factor of 2.0, 1.5, or 1.0 fold to the level of the RISKMARKER or INJURYMARKER transcript in the reference cell population. In various embodiments, a RISKMARKER or INJURYMARKER sequence in a test cell population can be considered altered in levels of expression if its expression level varies from the reference cell population by more than 1.0, 1.5, 2.0 or more fold from the expression level of the corresponding RISKMARKER or INJURYMARKER sequence in the reference cell population.

Alternatively, the absolute expression level matrix of the 8 RISKMARKER and/or 10 INJURYMARKER fragments in a test cell can be transformed using the principal component eigenvectors described above, or similar eigenvalues generated from parallel dosed members of the training set as internal controls. The expression eigenvalues for the test cell can then be compared to the training set eigenvalues described herein, or a parallel-run training set, if any.

The RISKMARKER expression level combination is considered similar to Low Risk idiosyncratic NSAIDS (several of which have been withdrawn), if the test drug's expression profile is within the 95% confidence interval (CI) of the centroid of that risk class. *See* Table 4. The test drug is considered Very Low Risk idiosyncratic if the transformed expression profile falls within the 95% CI of the centroid of that class. The test drug is considered Overdose Risk if the expression profile falls within the 95% CI of the centroid of that class. If the compound fails to associate with any of these compounds it will be categorized as unclassifiable.

Similarly, the INJURYMARKER expression level combination is considered indicative of hepatocellular damage induced by idiosyncratic NSAIDS, if the test drug's expression profile is within the 95% confidence interval (CI) of the centroid of that injury type. *See* Table 6. The test drug is considered to induce idiosyncratic cholestasis if the transformed expression profile falls within the 95% CI of the centroid of that injury type. The test drug is considered to induce elevated transaminase level if the expression profile falls within the 95% CI of the centroid of that class. If the compound fails to associate with any of these compounds it will be categorized as unclassifiable.

If desired, comparison of differentially expressed sequences between a test cell population and a reference cell population can be done with respect to a control nucleic acid whose expression is independent of the parameter or condition being measured. Expression levels of the control nucleic acid in the test and reference nucleic acid can be used to normalize signal levels in the compared populations.

In some embodiments, the test cell population is compared to multiple reference cell populations. Each of the multiple reference populations may differ in the known parameter. Thus, a test cell population may be compared to a first reference cell population known to have been exposed to a hepatotoxic agent, as well as a second reference population known have not been exposed to a hepatotoxic agent.

The test cell population that is exposed to, *i.e.*, contacted with, the test hepatotoxic agent can be any number of cells, *i.e.*, one or more cells, and can be provided *in vitro*, *in vivo*, or *ex vivo*.

In other embodiments, the test cell population can be divided into two or more subpopulations. The subpopulations can be created by dividing the first population of cells to create as identical a subpopulation as possible. This will be suitable, in, for example, *in vitro* or *ex vivo* screening methods. In some embodiments, various sub populations can be exposed

to a control agent, and/or a test agent, multiple test agents, or, *e.g.*, varying dosages of one or multiple test agents administered together, or in various combinations.

Preferably, cells in the reference cell population are derived from a tissue type as similar as possible to test cell, *e.g.*, liver tissue. In some embodiments, the control cell is derived from the same subject as the test cell, *e.g.*, from a region proximal to the region of origin of the test cell. In other embodiments, the reference cell population is derived from a plurality of cells. For example, the reference cell population can be a database of expression patterns from previously tested cells for which one of the herein-described parameters or conditions (hepatotoxic agent expression status is known.

10 The test agent can be a compound not previously described or can be a previously known compound but which is not known to be a hepatotoxic agent.

The subject is preferably a mammal. The mammal can be, *e.g.*, a human, non-human primate, mouse, rat, dog, cat, horse, or cow.

SCREENING FOR TOXIC AGENTS

15 In one aspect, the invention provides a method of identifying toxic agents, *e.g.*, hepatotoxic agents. The hepatotoxic agent can be identified by providing a cell population that includes cells capable of expressing one or more nucleic acid sequences homologous to those of RISKMARKER 1-8 or INJURYMARKER 1-10. The sequences need not be identical to sequences including RISKMARKER or INJURYMARKER nucleic acid sequences, as long as the sequence is sufficiently similar that specific hybridization can be detected. Preferably, the cell includes sequences that are identical, or nearly identical to those identifying the RISKMARKER or INJURYMARKER nucleic acids described herein.

Expression of the nucleic acid sequences in the test cell population is then compared to the expression of the nucleic acid sequences in a reference cell population, which is a cell population that has not been exposed to the test agent, or, in some embodiments, a cell population exposed the test agent. Comparison can be performed on test and reference samples measured concurrently or at temporally distinct times. An example of the latter is the use of compiled expression information, *e.g.*, a sequence database, which assembles information about expression levels of known sequences following administration of various agents. For example, alteration of expression levels following administration of test agent can be compared to the expression changes observed in the nucleic acid sequences following administration of a control agent, *e.g.* a NSAID such as ketoprofen.

An alteration in expression of the nucleic acid sequence in the test cell population compared to the expression of the nucleic acid sequence in the reference cell population that has not been exposed to the test agent indicates the test agent is a hepatotoxic agent.

The invention also includes a hepatotoxic agent identified according to this screening
5 method.

In some embodiments of the method of the invention, the test agent is an idiosyncratic hepatotoxic agent, *e.g.* a NSAID, and the reference agent is also a NSAID. As described above, RISKMARKER (*e.g.* RISKMARKERS 1-8) expression level patterns can be used to predict the level of hepatotoxicity risk (*i.e.* low, very low, or overdose) associated with a given
10 test agent, *e.g.* a NSAID. In one embodiment, the reference NSAID (*i.e.* used with the reference cell population) is a NSAID classified as having a low risk of hepatotoxicity. The test agent is then identified as having a low risk of hepatotoxicity if no qualitative difference in expression levels is identified in comparison to expression levels in the reference population exposed to a low risk NSAID. In certain embodiments, the low risk NSAID is Benoxaprofen,
15 Bromfenac, Diclofenac, Phenylbutazone, or Sulindac. In another embodiment, the reference NSAID is a NSAID classified as having a very low risk of hepatotoxicity. The test agent is then identified as having a very low risk of hepatotoxicity if no qualitative difference in expression levels is identified in comparison to expression levels in the reference population exposed to a very low risk NSAID. In certain embodiments, the very low risk NSAID is Etodolac,
20 Fenoprofen, Flurbiprofen, Ibuprofen, Indomethacin, Ketoprofen, Meclofenamate, Mefenamic Acid, Nabumetone, Naproxen, Oxaprozin, Piroxicam, Suprofen, Tenoxicam, Tolmentin, or Zomepirac. In still another embodiment, the reference NSAID is a NSAID classified as having an overdose risk of hepatotoxicity. The test agent is then identified as having an overdose risk of hepatotoxicity if no qualitative difference in expression levels is identified in
25 comparison to expression levels in the reference population exposed to an overdose risk NSAID. In certain embodiments, the overdose risk NSAID is Acetaminophen, Acetylsalicylic acid, or Phenacetin. In some embodiments, the difference in expression levels is determined by comparing expression transformation eigenvectors (for risk class) for the test cell and reference cell populations, as described above.

30 As also described above, INJURYMARKER (*e.g.* INJURYMARKERS 1-10) expression level patterns can be used to predict the type of hepatotoxicity injury (*i.e.* hepatocellular damage, cholestasis, or elevated transaminase level) associated with a given test agent, *e.g.* a NSAID. In some embodiments, the reference NSAID is a NSAID classified as

inducing hepatocellular damage. The test agent is then identified as likely to induce hepatocellular damage if no qualitative difference in expression levels is identified in comparison to expression levels in the reference population exposed to a NSAID which induces hepatocellular damage. In certain embodiments, the hepatocellular damage inducing NSAID is Acetaminophen, Flurbiprofen, or Ketoprofen. In another embodiment, the reference NSAID is a NSAID classified as inducing cholestasis. The test agent is then identified as likely to induce cholestasis if no qualitative difference in expression levels is identified in comparison to expression levels in the reference population exposed to a NSAID which induces cholestasis. In certain embodiments, the cholestasis-inducing NSAID is Benoxaprofen, Nabumetone, or Sulindac. In yet another embodiment, the reference NSAID is a NSAID classified as inducing elevated transaminase level. The test agent is then identified as likely to induce elevated transaminase level if no qualitative difference in expression levels is identified as compared to expression levels in the reference population exposed to a NSAID which induces elevated transaminase levels. In certain embodiments, the elevated transaminase level inducing NSAID is Zomepirac, Mefenamic acid, or Tenoxicam. In some embodiments, the difference in expression levels is determined by comparing expression transformation eigenvectors for said test cell and reference cell populations, as described above.

ASSESSING TOXICITY OF A TOXIC AGENT IN A SUBJECT

The differentially expressed RISKMARKER or INJURYMARKER sequences identified herein also allow for the hepatotoxicity of a hepatotoxic agent to be determined or monitored. In this method, a test cell population from a subject is exposed to a test agent, *i.e.* a hepatotoxic agent. If desired, test cell populations can be taken from the subject at various time points before, during, or after exposure to the test agent. Expression of one or more of the RISKMARKER or INJURYMARKER sequences in the cell population is then measured and compared to a reference cell population which includes cells whose hepatotoxic agent expression status is known. Preferably, the reference cells not been exposed to the test agent.

If the reference cell population contains no cells exposed to the treatment, a similarity in expression between RISKMARKER or INJURYMARKER sequences in the test cell population and the reference cell population indicates that the treatment is non-hepatotoxic. However, a difference in expression between RISKMARKER or INJURYMARKER sequences in the test population and this reference cell population indicates the treatment is hepatotoxic.

By "hepatotoxicity" is meant that the agent is damaging or destructive to liver when administered to a subject leads to liver damage.

As described in detail above, RISKMARKER expression patterns can be used to predict the level of hepatotoxicity risk (*e.g.* low risk, very low risk, overdose risk) associated with a test agent or drug, by comparison to RISKMARKER expression levels for reference drugs, *e.g.* NSAIDs, with a given classification of risk (*e.g.* very low risk). Similarly, INJURYMARKER expression patterns can be used to predict the type of hepatotoxicity damage (*e.g.* hepatocellular damage, cholestasis, elevated transaminase level) associated with a test agent or drug, by comparison to INJURYMARKER expression levels for reference drugs, *e.g.* NSAIDs, which induce a given type of hepatotoxic damage (*e.g.* cholestasis).

RISKMARKER NUCLEIC ACIDS

Also provided in the invention are novel nucleic acid comprising a nucleic acid sequence selected from the group consisting of RISKMARKER 1, and RISKMARKERS 6-8, or their complements, as well as vectors and cells including these nucleic acids.

Thus, one aspect of the invention pertains to isolated RISKMARKER nucleic acid molecules that encode RISKMARKER proteins or biologically active portions thereof. Also included are nucleic acid fragments sufficient for use as hybridization probes to identify RISKMARKER-encoding nucleic acids (*e.g.*, RISKMARKER mRNA) and fragments for use as polymerase chain reaction (PCR) primers for the amplification or mutation of RISKMARKER nucleic acid molecules. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (*e.g.*, cDNA or genomic DNA), RNA molecules (*e.g.*, mRNA), analogs of the DNA or RNA generated using nucleotide analogs, and derivatives, fragments and homologs thereof. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

"Probes" refer to nucleic acid sequences of variable length, preferably between at least about 10 nucleotides (nt) or as many as about, *e.g.*, 6,000 nt, depending on use. Probes are used in the detection of identical, similar, or complementary nucleic acid sequences. Longer length probes are usually obtained from a natural or recombinant source, are highly specific and much slower to hybridize than oligomers. Probes may be single- or double-stranded and designed to have specificity in PCR, membrane-based hybridization technologies, or ELISA-like technologies.

An "isolated" nucleic acid molecule is one that is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Examples of isolated nucleic acid molecules include, but are not limited to, recombinant DNA molecules contained in a vector, recombinant DNA molecules maintained in a heterologous host cell, partially or substantially purified nucleic acid molecules, and synthetic DNA or RNA molecules. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (*i.e.*, sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated RISKMARKER nucleic acid molecule can contain less than about 50 kb, 25 kb, 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material or culture medium when produced by recombinant techniques, or of chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, *e.g.*, a nucleic acid molecule having the nucleotide sequence of any of RISKMARKER 1, or RISKMARKER 6-8, or a complement of any of these nucleotide sequences, can be isolated using standard molecular biology techniques and the sequence information provided herein. Using all or a portion of these nucleic acid sequences as a hybridization probe, RISKMARKER or INJURYMARKER nucleic acid sequences can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook *et al.*, eds., Molecular Cloning: A Laboratory Manual 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989; and Ausubel, *et al.*, eds., Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, 1993.)

A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to RISKMARKER nucleotide sequences can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

As used herein, the term "oligonucleotide" refers to a series of linked nucleotide residues, which oligonucleotide has a sufficient number of nucleotide bases to be used in a PCR reaction. A short oligonucleotide sequence may be based on, or designed from, a genomic or cDNA sequence and is used to amplify, confirm, or reveal the presence of an identical, similar or complementary DNA or RNA in a particular cell or tissue.

Oligonucleotides comprise portions of a nucleic acid sequence having at least about 10 nt and as many as 50 nt, preferably about 15 nt to 30 nt. They may be chemically synthesized and may be used as probes.

In another embodiment, an isolated nucleic acid molecule of the invention comprises a
5 nucleic acid molecule that is a complement of the nucleotide sequence shown in
RISKMARKER 1, or RISKMARKER 6-8. In another embodiment, an isolated nucleic acid
molecule of the invention comprises a nucleic acid molecule that is a complement of the
nucleotide sequence shown in any of these sequences, or a portion of any of these nucleotide
sequences. A nucleic acid molecule that is complementary to the nucleotide sequence shown
10 in RISKMARKER 1, or RISKMARKER 6-8 is one that is sufficiently complementary to the
nucleotide sequence shown, such that it can hydrogen bond with little or no mismatches to the
nucleotide sequences shown, thereby forming a stable duplex.

As used herein, the term "complementary" refers to Watson-Crick or Hoogsteen base
pairing between nucleotides units of a nucleic acid molecule, and the term "binding" means
15 the physical or chemical interaction between two polypeptides or compounds or associated
polypeptides or compounds or combinations thereof. Binding includes ionic, non-ionic, Von
der Waals, hydrophobic interactions, etc. A physical interaction can be either direct or
indirect. Indirect interactions may be through or due to the effects of another polypeptide or
compound. Direct binding refers to interactions that do not take place through, or due to, the
20 effect of another polypeptide or compound, but instead are without other substantial chemical
intermediates.

Moreover, the nucleic acid molecule of the invention can comprise only a portion of
the nucleic acid sequence of RISKMARKER 1, or RISKMARKER 6-8, *e.g.*, a fragment that
can be used as a probe or primer or a fragment encoding a biologically active portion of
25 RISKMARKER. Fragments provided herein are defined as sequences of at least 6
(contiguous) nucleic acids or at least 4 (contiguous) amino acids, a length sufficient to allow
for specific hybridization in the case of nucleic acids or for specific recognition of an epitope
in the case of amino acids, respectively, and are at most some portion less than a full length
sequence. Fragments may be derived from any contiguous portion of a nucleic acid or amino
30 acid sequence of choice. Derivatives are nucleic acid sequences or amino acid sequences
formed from the native compounds either directly or by modification or partial substitution.
Analogous are nucleic acid sequences or amino acid sequences that have a structure similar to,
but not identical to, the native compound but differs from it in respect to certain components

or side chains. Analogs may be synthetic or from a different evolutionary origin and may have a similar or opposite metabolic activity compared to wild type.

Derivatives and analogs may be full length or other than full length, if the derivative or analog contains a modified nucleic acid or amino acid, as described below. Derivatives or analogs of the nucleic acids or proteins of the invention include, but are not limited to, molecules comprising regions that are substantially homologous to the nucleic acids or proteins of the invention, in various embodiments, by at least about 45%, 50%, 70%, 80%, 95%, 98%, or even 99% identity (with a preferred identity of 80-99%) over a nucleic acid or amino acid sequence of identical size or when compared to an aligned sequence in which the alignment is done by a computer homology program known in the art, or whose encoding nucleic acid is capable of hybridizing to the complement of a sequence encoding the aforementioned proteins under stringent, moderately stringent, or low stringent conditions. See *e.g.* Ausubel, *et al.*, Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, 1993, and below. An exemplary program is the Gap program (Wisconsin Sequence Analysis Package, Version 8 for UNIX, Genetics Computer Group, University Research Park, Madison, WI) using the default settings, which uses the algorithm of Smith and Waterman (Adv. Appl. Math., 1981, 2: 482-489, which is incorporated herein by reference in its entirety).

A "homologous nucleic acid sequence" or "homologous amino acid sequence," or variations thereof, refer to sequences characterized by a homology at the nucleotide level or amino acid level as discussed above. Homologous nucleotide sequences encode those sequences coding for isoforms of a RISKMARKER polypeptide. Isoforms can be expressed in different tissues of the same organism as a result of, for example, alternative splicing of RNA. Alternatively, isoforms can be encoded by different genes. In the present invention, homologous nucleotide sequences include nucleotide sequences encoding for a RISKMARKER polypeptide of species other than humans, including, but not limited to, mammals, and thus can include, *e.g.*, mouse, rat, rabbit, dog, cat, cow, horse, and other organisms. Homologous nucleotide sequences also include, but are not limited to, naturally occurring allelic variations and mutations of the nucleotide sequences set forth herein. A homologous nucleotide sequence does not, however, include the nucleotide sequence encoding a human RISKMARKER protein. Homologous nucleic acid sequences include those nucleic acid sequences that encode conservative amino acid substitutions (see below) in a RISKMARKER polypeptide, as well as a polypeptide having a RISKMARKER activity. A

homologous amino acid sequence does not encode the amino acid sequence of a human RISKMARKER polypeptide.

The nucleotide sequence determined from the cloning of human RISKMARKER genes allows for the generation of probes and primers designed for use in identifying and/or cloning
5 RISKMARKER homologues in other cell types, *e.g.*, from other tissues, as well as RISKMARKER homologues from other mammals. The probe/primer typically comprises a substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 25, 50, 100, 150, 200, 250, 300, 350 or 400 consecutive sense strand nucleotide sequence of a nucleic acid
10 comprising a RISKMARKER sequence, or an anti-sense strand nucleotide sequence of a nucleic acid comprising a RISKMARKER sequence, or of a naturally occurring mutant of these sequences.

Probes based on human RISKMARKER nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In various
15 embodiments, the probe further comprises a label group attached thereto, *e.g.*, the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress a RISKMARKER protein, such as by measuring a level of a RISKMARKER-encoding nucleic acid in a sample of cells from a subject *e.g.*, detecting
20 RISKMARKER mRNA levels or determining whether a genomic RISKMARKER gene has been mutated or deleted.

"A polypeptide having a biologically active portion of RISKMARKER" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular
25 biological assay, with or without dose dependency. A nucleic acid fragment encoding a "biologically active portion of RISKMARKER" can be prepared by isolating a portion of RISKMARKER 1, or RISKMARKER 6-8, that encodes a polypeptide having a RISKMARKER biological activity, expressing the encoded portion of RISKMARKER protein (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of
30 RISKMARKER. For example, a nucleic acid fragment encoding a biologically active portion of a RISKMARKER polypeptide can optionally include an ATP-binding domain. In another embodiment, a nucleic acid fragment encoding a biologically active portion of RISKMARKER includes one or more regions.

RISKMARKER AND INJURYMARKER VARIANTS

The invention further encompasses nucleic acid molecules that differ from the disclosed or referenced RISKMARKER or INJURYMARKER nucleotide sequences due to degeneracy of the genetic code. These nucleic acids thus encode the same RISKMARKER or INJURYMARKER protein as that encoded by nucleotide sequence comprising a RISKMARKER or INJURYMARKER nucleic acid as shown in, *e.g.*, RISKMARKER 1-8 INJURYMARKER 1-10.

In addition to the rat RISKMARKER or INJURYMARKER nucleotide sequence shown in RISKMARKER or INJURYMARKER 1, and RISKMARKER or INJURYMARKER 6-8, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of a RISKMARKER or INJURYMARKER polypeptide may exist within a population (*e.g.*, the human population). Such genetic polymorphism in the RISKMARKER or INJURYMARKER gene may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a RISKMARKER or INJURYMARKER protein, preferably a mammalian RISKMARKER or INJURYMARKER protein. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of the RISKMARKER or INJURYMARKER gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in RISKMARKER or INJURYMARKER that are the result of natural allelic variation and that do not alter the functional activity of RISKMARKER or INJURYMARKER are intended to be within the scope of the invention.

Moreover, nucleic acid molecules encoding RISKMARKER or INJURYMARKER proteins from other species, and thus that have a nucleotide sequence that differs from the human sequence of RISKMARKER OR INJURYMARKER, are intended to be within the scope of the invention. Nucleic acid molecules corresponding to natural allelic variants and homologues of the RISKMARKER or INJURYMARKER DNAs of the invention can be isolated based on their homology to the human RISKMARKER or INJURYMARKER nucleic acids disclosed herein using the human cDNAs, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. For example, a soluble human RISKMARKER or INJURYMARKER DNA can be isolated based on its homology to human membrane-bound RISKMARKER or INJURYMARKER. Likewise, a membrane-bound human RISKMARKER or INJURYMARKER DNA can be isolated based on its homology to soluble human RISKMARKER or INJURYMARKER.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 6 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising the nucleotide sequence of RISKMARKER 1, or RISKMARKER 6-8. In another embodiment, the nucleic acid is at least 10, 25, 50, 100, 250
5 or 500 nucleotides in length. In another embodiment, an isolated nucleic acid molecule of the invention hybridizes to the coding region. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other.

10 Homologs (*i.e.*, nucleic acids encoding RISKMARKER or INJURYMARKER proteins derived from species other than human) or other related sequences (*e.g.*, paralogs) can be obtained by low, moderate or high stringency hybridization with all or a portion of the particular human sequence as a probe using methods well known in the art for nucleic acid hybridization and cloning.

15 As used herein, the phrase "stringent hybridization conditions" refers to conditions under which a probe, primer or oligonucleotide will hybridize to its target sequence, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures than shorter sequences. Generally, stringent conditions are selected to be about 5°C lower than the
20 thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the probes complementary to the target sequence hybridize to the target sequence at equilibrium. Since the target sequences are generally present at excess, at T_m, 50% of the probes are occupied at equilibrium. Typically, stringent conditions will be those in
25 which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes, primers or oligonucleotides (*e.g.*, 10 nt to 50 nt) and at least about 60°C for longer probes, primers and oligonucleotides. Stringent conditions may also be achieved with the addition of destabilizing agents, such as formamide.

30 Stringent conditions are known to those skilled in the art and can be found in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Preferably, the conditions are such that sequences at least about 65%, 70%, 75%, 85%, 90%, 95%, 98%, or 99% homologous to each other typically remain hybridized to each other. A non-limiting example of stringent hybridization conditions is hybridization in a high salt buffer comprising

6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 mg/ml denatured salmon sperm DNA at 65°C. This hybridization is followed by one or more washes in 0.2X SSC, 0.01% BSA at 50°C. An isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of RISKMARKER 1, or
5 RISKMARKER 6-8 corresponds to a naturally occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

In a second embodiment, a nucleic acid sequence that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of RISKMARKER 1, or RISKMARKER
10 6-8, or fragments, analogs or derivatives thereof, under conditions of moderate stringency is provided. A non-limiting example of moderate stringency hybridization conditions are hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100 mg/ml denatured salmon sperm DNA at 55°C, followed by one or more washes in 1X SSC, 0.1% SDS at 37°C. Other conditions of moderate stringency that may be used are well known in the art. See, *e.g.*,
15 Ausubel *et al.* (eds.), 1993, Current Protocols in Molecular Biology, John Wiley & Sons, NY, and Kriegler, 1990, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY.

In a third embodiment, a nucleic acid that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of RISKMARKER 1, or RISKMARKER 6-8, or fragments, analogs or derivatives thereof, under conditions of low stringency, is provided. A
20 non-limiting example of low stringency hybridization conditions are hybridization in 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 mg/ml denatured salmon sperm DNA, 10% (wt/vol) dextran sulfate at 40°C, followed by one or more washes in 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS at 50°C. Other conditions of low stringency that may be used are well known in
25 the art (*e.g.*, as employed for cross-species hybridizations). See, *e.g.*, Ausubel *et al.* (eds.), 1993, Current Protocols in Molecular Biology, John Wiley & Sons, NY, and Kriegler, 1990, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY; Shilo *et al.*, 1981, *Proc Natl Acad Sci USA* 78: 6789-6792.

CONSERVATIVE MUTATIONS

30 In addition to naturally-occurring allelic variants of the RISKMARKER sequence that may exist in the population, the skilled artisan will further appreciate that changes can be introduced into an RISKMARKER nucleic acid or directly into an RISKMARKER

polypeptide sequence without altering the functional ability of the RISKMARKER protein. In some embodiments, the nucleotide sequence of RISKMARKER 1, or RISKMARKER 6-8 will be altered, thereby leading to changes in the amino acid sequence of the encoded RISKMARKER protein. For example, nucleotide substitutions that result in amino acid
5 substitutions at various "non-essential" amino acid residues can be made in the sequence of RISKMARKER 1, or RISKMARKER 6-8. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of RISKMARKER without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are conserved among the RISKMARKER
10 proteins of the present invention, are predicted to be particularly unamenable to alteration.

In addition, amino acid residues that are conserved among family members of the RISKMARKER proteins of the present invention, are also predicted to be particularly unamenable to alteration. As such, these conserved domains are not likely to be amenable to mutation. Other amino acid residues, however, (*e.g.*, those that are not conserved or only
15 semi-conserved among members of the RISKMARKER proteins) may not be essential for activity and thus are likely to be amenable to alteration.

Another aspect of the invention pertains to nucleic acid molecules encoding RISKMARKER proteins that contain changes in amino acid residues that are not essential for activity. Such RISKMARKER proteins differ in amino acid sequence from the amino acid
20 sequences of polypeptides encoded by nucleic acids containing RISKMARKER 1, or RISKMARKER 6-8, yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 45% homologous, more preferably 60%, and still more preferably at least about 70%, 80%, 90%, 95%, 98%, and most preferably at least
25 about 99% homologous to the amino acid sequence of the amino acid sequences of polypeptides encoded by nucleic acids comprising RISKMARKER 1, or RISKMARKER 6-8.

An isolated nucleic acid molecule encoding a RISKMARKER protein homologous to can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of a nucleic acid comprising RISKMARKER 1, or RISKMARKER
30 6-8, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein.

Mutations can be introduced into a nucleic acid comprising RISKMARKER 1, or RISKMARKER 6-8 by standard techniques, such as site-directed mutagenesis and

PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have
5 been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side
10 chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in RISKMARKER is replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a RISKMARKER coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for RISKMARKER biological activity
15 to identify mutants that retain activity. Following mutagenesis of the nucleic acid, the encoded protein can be expressed by any recombinant technology known in the art and the activity of the protein can be determined.

In one embodiment, a mutant RISKMARKER protein can be assayed for (1) the ability to form protein:protein interactions with other RISKMARKER proteins, other cell-surface
20 proteins, or biologically active portions thereof, (2) complex formation between a mutant RISKMARKER protein and a RISKMARKER ligand; (3) the ability of a mutant RISKMARKER protein to bind to an intracellular target protein or biologically active portion thereof; (*e.g.*, avidin proteins); (4) the ability to bind ATP; or (5) the ability to specifically bind a RISKMARKER protein antibody.

25 In other embodiment, the fragment of the complementary polynucleotide sequence described in claim 1 wherein the fragment of the complementary polynucleotide sequence hybridizes to the first sequence.

In other specific embodiments, the nucleic acid is RNA or DNA. The fragment or the fragment of the complementary polynucleotide sequence described in claim 38, wherein the
30 fragment is between about 10 and about 100 nucleotides in length, *e.g.*, between about 10 and about 90 nucleotides in length, or about 10 and about 75 nucleotides in length, about 10 and about 50 bases in length, about 10 and about 40 bases in length, or about 15 and about 30 bases in length.

Another aspect of the invention pertains to isolated antisense nucleic acid molecules that are hybridizable to or complementary to the nucleic acid molecule comprising the nucleotide sequence of a RISKMARKER or INJURYMARKER sequence or fragments, analogs or derivatives thereof. An "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. In specific aspects, antisense nucleic acid molecules are provided that comprise a sequence complementary to at least about 10, 25, 50, 100, 250 or 500 nucleotides or an entire RISKMARKER or INJURYMARKER coding strand, or to only a portion thereof. Nucleic acid molecules encoding fragments, homologs, derivatives and analogs of a RISKMARKER or INJURYMARKER protein, or antisense nucleic acids complementary to a nucleic acid comprising a RISKMARKER or INJURYMARKER nucleic acid sequence are additionally provided.

In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding RISKMARKER or INJURYMARKER. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues. In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding RISKMARKER. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences encoding RISKMARKER or INJURYMARKER disclosed herein, antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of RISKMARKER or INJURYMARKER mRNA, but more preferably is an oligonucleotide that is antisense to only a portion of the coding or noncoding region of RISKMARKER or INJURYMARKER mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of RISKMARKER or INJURYMARKER mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis or enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally

occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used.

5 Examples of modified nucleotides that can be used to generate the antisense nucleic acid include: 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine,
10 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil,
15 uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest,
20 described further in the following subsection).

 The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a RISKMARKER or INJURYMARKER protein to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization
25 can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and
30 then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient

intracellular concentrations of antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an
5 α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual α -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res* 15: 6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res* 15: 6131-6148) or a chimeric RNA -DNA analogue (Inoue *et al.*
10 (1987) *FEBS Lett* 215: 327-330).

RIBOZYMES AND PNA MOIETIES

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity that are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a
15 complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave RISKMARKER or INJURYMARKER mRNA transcripts to thereby inhibit translation of RISKMARKER or INJURYMARKER mRNA. A ribozyme having specificity for a RISKMARKER or INJURYMARKER -encoding nucleic acid can be designed based upon the
20 nucleotide sequence of a RISKMARKER or INJURYMARKER DNA disclosed herein. For example, a derivative of a Tetrahymena L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a RISKMARKER or INJURYMARKER-encoding mRNA. See, *e.g.*, Cech *et al.* U.S. Pat. No. 4,987,071; and Cech *et al.* U.S. Pat. No. 5,116,742. Alternatively,
25 RISKMARKER or INJURYMARKER mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel *et al.*, (1993) *Science* 261:1411-1418.

Alternatively, RISKMARKER or INJURYMARKER gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of a
30 RISKMARKER or INJURYMARKER nucleic acid (*e.g.*, the RISKMARKER or INJURYMARKER promoter and/or enhancers) to form triple helical structures that prevent transcription of the RISKMARKER or INJURYMARKER gene in target cells. See generally,

Helene. (1991) *Anticancer Drug Des.* 6: 569-84; Helene. *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher (1992) *Bioassays* 14: 807-15.

In various embodiments, the nucleic acids of RISKMARKER or INJURYMARKER can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids (see Hyrup *et al.* (1996) *Bioorg Med Chem* 4: 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup *et al.* (1996) above; Perry-O'Keefe *et al.* (1996) *PNAS* 93: 14670-675.

PNAs of RISKMARKER or INJURYMARKER can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, *e.g.*, inducing transcription or translation arrest or inhibiting replication. PNAs of RISKMARKER or INJURYMARKER can also be used, *e.g.*, in the analysis of single base pair mutations in a gene by, *e.g.*, PNA directed PCR clamping; as artificial restriction enzymes when used in combination with other enzymes, *e.g.*, S1 nucleases (Hyrup B. (1996) above); or as probes or primers for DNA sequence and hybridization (Hyrup *et al.* (1996), above; Perry-O'Keefe (1996), above).

In another embodiment, PNAs of RISKMARKER or INJURYMARKER can be modified, *e.g.*, to enhance their stability or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras of RISKMARKER or INJURYMARKER can be generated that may combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, *e.g.*, RNase H and DNA polymerases, to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup (1996) above). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup (1996) above and Finn *et al.* (1996) *Nucl Acids Res* 24: 3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry, and modified nucleoside analogs, *e.g.*,

5'- (4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used between the PNA and the 5' end of DNA (Mag *et al.* (1989) *Nucl Acid Res* 17: 5973-88). PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn *et al.* (1996) above). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment. See, Petersen *et al.* (1975) *Bioorg Med Chem Lett* 5: 1119-11124.

In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger *et al.*, 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556; Lemaitre *et al.*, 1987, *Proc. Natl. Acad. Sci.* 84:648-652; PCT Publication No. W088/09810) or the blood-brain barrier (see, *e.g.*, PCT Publication No. W089/10134). In addition, oligonucleotides can be modified with hybridization triggered cleavage agents (See, *e.g.*, Krol *et al.*, 1988, *BioTechniques* 6:958-976) or intercalating agents. (See, *e.g.*, Zon, 1988, *Pharm. Res.* 5: 539-549). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, a hybridization triggered cross-linking agent, a transport agent, a hybridization-triggered cleavage agent, etc.

RISKMARKER AND INJURYMARKER POLYPEPTIDES

One aspect of the invention pertains to isolated RISKMARKER or INJURYMARKER proteins, and biologically active portions thereof, or derivatives, fragments, analogs or homologs thereof. Also provided are polypeptide fragments suitable for use as immunogens to raise anti-RISKMARKER or INJURYMARKER antibodies, *e.g.* antibodies against RISKMARKER 1, or RISKMARKER 6-8. In one embodiment, native RISKMARKER or INJURYMARKER proteins can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, RISKMARKER or INJURYMARKER proteins are produced by recombinant DNA techniques. Alternative to recombinant expression, a RISKMARKER or INJURYMARKER protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the RISKMARKER or INJURYMARKER protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language

"substantially free of cellular material" includes preparations of RISKMARKER or INJURYMARKER protein in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of RISKMARKER or INJURYMARKER protein having less than about 30% (by dry weight) of non-RISKMARKER or INJURYMARKER protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-RISKMARKER or INJURYMARKER protein, still more preferably less than about 10% of non-RISKMARKER or INJURYMARKER protein, and most preferably less than about 5% non-RISKMARKER or INJURYMARKER protein. When the RISKMARKER or INJURYMARKER protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of RISKMARKER or INJURYMARKER protein in which the protein is separated from chemical precursors or other chemicals that are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of RISKMARKER or INJURYMARKER protein having less than about 30% (by dry weight) of chemical precursors or non-RISKMARKER or INJURYMARKER chemicals, more preferably less than about 20% chemical precursors or non-RISKMARKER or INJURYMARKER chemicals, still more preferably less than about 10% chemical precursors or non-RISKMARKER or INJURYMARKER chemicals, and most preferably less than about 5% chemical precursors or non-RISKMARKER or INJURYMARKER chemicals.

Biologically active portions of a RISKMARKER or INJURYMARKER protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the RISKMARKER or INJURYMARKER protein, *e.g.*, the amino acid sequence encoded by a nucleic acid comprising RISKMARKER or INJURYMARKER 1-20 that include fewer amino acids than the full length RISKMARKER or INJURYMARKER proteins, and exhibit at least one activity of a RISKMARKER or INJURYMARKER protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the RISKMARKER or INJURYMARKER protein. A biologically active portion of a

RISKMARKER or INJURYMARKER protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length.

A biologically active portion of a RISKMARKER or INJURYMARKER protein of the present invention may contain at least one of the above-identified domains conserved between the RISKMARKER or INJURYMARKER proteins. An alternative biologically active portion of a RISKMARKER or INJURYMARKER protein may contain at least two of the above-identified domains. Another biologically active portion of a RISKMARKER or INJURYMARKER protein may contain at least three of the above-identified domains. Yet another biologically active portion of a RISKMARKER or INJURYMARKER protein of the present invention may contain at least four of the above-identified domains.

Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native RISKMARKER or INJURYMARKER protein.

In some embodiments, the RISKMARKER or INJURYMARKER protein is substantially homologous to one of these RISKMARKER or INJURYMARKER proteins and retains its the functional activity, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail below.

In specific embodiments, the invention includes an isolated polypeptide comprising an amino acid sequence that is 80% or more identical to the sequence of a polypeptide whose expression is modulated in a mammal to which hepatotoxic agent is administered.

DETERMINING HOMOLOGY BETWEEN TWO OR MORE SEQUENCES

To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are homologous at that position (*i.e.*, as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity").

The nucleic acid sequence homology may be determined as the degree of identity between two sequences. The homology may be determined using computer programs known in the art, such as GAP software provided in the GCG program package. See *Needleman and Wunsch* 1970 *J Mol Biol* 48: 443-453. Using GCG GAP software with the following settings
5 for nucleic acid sequence comparison: GAP creation penalty of 5.0 and GAP extension penalty of 0.3, the coding region of the analogous nucleic acid sequences referred to above exhibits a degree of identity preferably of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99%, with the CDS (encoding) part of a DNA sequence comprising RISKMARKER 1, or RISKMARKER 6-8..

10 The term "sequence identity" refers to the degree to which two polynucleotide or polypeptide sequences are identical on a residue-by-residue basis over a particular region of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over that region of comparison, determining the number of positions at which the identical nucleic acid base (*e.g.*, A, T, C, G, U, or I, in the case of
15 nucleic acids) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the region of comparison (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The term "substantial identity" as used herein denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 80
20 percent sequence identity, preferably at least 85 percent identity and often 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison region.

CHIMERIC AND FUSION PROTEINS

The invention also provides RISKMARKER chimeric or fusion proteins. As used
25 herein, an RISKMARKER "chimeric protein" or "fusion protein" comprises an RISKMARKER polypeptide operatively linked to a non-RISKMARKER polypeptide. A "RISKMARKER polypeptide" refers to a polypeptide having an amino acid sequence corresponding to RISKMARKER, whereas a "non-RISKMARKER polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein that is not substantially
30 homologous to the RISKMARKER protein, *e.g.*, a protein that is different from the RISKMARKER protein and that is derived from the same or a different organism. Within an RISKMARKER fusion protein the RISKMARKER polypeptide can correspond to all or a

portion of an RISKMARKER protein. In one embodiment, an RISKMARKER fusion protein comprises at least one biologically active portion of an RISKMARKER protein. In another embodiment, an RISKMARKER fusion protein comprises at least two biologically active portions of an RISKMARKER protein. In yet another embodiment, an RISKMARKER fusion protein comprises at least three biologically active portions of an RISKMARKER protein. Within the fusion protein, the term "operatively linked" is intended to indicate that the RISKMARKER polypeptide and the non-RISKMARKER polypeptide are fused in-frame to each other. The non-RISKMARKER polypeptide can be fused to the N-terminus or C-terminus of the RISKMARKER polypeptide.

For example, in one embodiment an RISKMARKER fusion protein comprises an RISKMARKER domain operably linked to the extracellular domain of a second protein. Such fusion proteins can be further utilized in screening assays for compounds which modulate RISKMARKER activity (such assays are described in detail below).

In yet another embodiment, the fusion protein is a GST-RISKMARKER fusion protein in which the RISKMARKER sequences are fused to the C-terminus of the GST (*i.e.*, glutathione S-transferase) sequences. Such fusion proteins can facilitate the purification of recombinant RISKMARKER, *e.g.* RISKMARKER 1, or RISKMARKER 6-8.

In another embodiment, the fusion protein is an RISKMARKER protein containing a heterologous signal sequence at its N-terminus. For example, a native RISKMARKER signal sequence can be removed and replaced with a signal sequence from another protein. In certain host cells (*e.g.*, mammalian host cells), expression and/or secretion of RISKMARKER can be increased through use of a heterologous signal sequence.

In yet another embodiment, the fusion protein is a RISKMARKER-immunoglobulin fusion protein in which the RISKMARKER sequences comprising one or more domains are fused to sequences derived from a member of the immunoglobulin protein family. The RISKMARKER-immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject to inhibit an interaction between a RISKMARKER ligand and a RISKMARKER protein on the surface of a cell, to thereby suppress RISKMARKER-mediated signal transduction *in vivo*. The RISKMARKER-immunoglobulin fusion proteins can be used to affect the bioavailability of an RISKMARKER cognate ligand. Inhibition of the RISKMARKER ligand/RISKMARKER interaction may be useful therapeutically for both the treatments of proliferative and differentiative disorders, as well as modulating (*e.g.* promoting or inhibiting) cell survival. Moreover, the RISKMARKER

-immunoglobulin fusion proteins of the invention can be used as immunogens to produce anti-RISKMARKER antibodies in a subject, to purify RISKMARKER ligands, and in screening assays to identify molecules that inhibit the interaction of RISKMARKER with a RISKMARKER ligand.

5 An RISKMARKER chimeric or fusion protein of the invention can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, *e.g.*, by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate,
10 alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers that give rise to complementary overhangs between two consecutive gene fragments that can subsequently be annealed and reamplified to generate a
15 chimeric gene sequence (see, for example, Ausubel *et al.* (eds.) Current Protocols in Molecular Biology, John Wiley & Sons, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). A RISKMARKER -encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the RISKMARKER protein.

20 **RISKMARKER AND INJURYMARKER AGONISTS AND ANTAGONISTS**

 The present invention also pertains to variants of the RISKMARKER or INJURYMARKER proteins that function as either RISKMARKER or INJURYMARKER agonists (mimetics) or as RISKMARKER or INJURYMARKER antagonists. Variants of the RISKMARKER or INJURYMARKER protein can be generated by mutagenesis, *e.g.*, discrete
25 point mutation or truncation of the RISKMARKER or INJURYMARKER protein. An agonist of the RISKMARKER or INJURYMARKER protein can retain substantially the same, or a subset of, the biological activities of the naturally occurring form of the RISKMARKER or INJURYMARKER protein. An antagonist of the RISKMARKER or INJURYMARKER protein can inhibit one or more of the activities of the naturally occurring form of the
30 RISKMARKER or INJURYMARKER protein by, for example, competitively binding to a downstream or upstream member of a cellular signaling cascade which includes the RISKMARKER or INJURYMARKER protein. Thus, specific biological effects can be

elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the RISKMARKER or INJURYMARKER proteins.

5 Variants of the RISKMARKER or INJURYMARKER protein that function as either RISKMARKER or INJURYMARKER agonists (mimetics) or as RISKMARKER or INJURYMARKER antagonists can be identified by screening combinatorial libraries of mutants, *e.g.*, truncation mutants, of the RISKMARKER or INJURYMARKER protein for RISKMARKER or INJURYMARKER protein agonist or antagonist activity. In one
10 embodiment, a variegated library of RISKMARKER or INJURYMARKER variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of RISKMARKER or INJURYMARKER variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential RISKMARKER
15 or INJURYMARKER sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display) containing the set of RISKMARKER or INJURYMARKER sequences therein. There are a variety of methods which can be used to produce libraries of potential RISKMARKER or INJURYMARKER variants from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can
20 be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential RISKMARKER or INJURYMARKER sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, *e.g.*, Narang (1983) *Tetrahedron* 39:3; Itakura *et al.* (1984) *Annu Rev Biochem* 53:323; Itakura *et al.* (1984) *Science* 198:1056; Ike *et al.* (1983) *Nucl Acid Res*
25 11:477.

POLYPEPTIDE LIBRARIES

In addition, libraries of fragments of the RISKMARKER or INJURYMARKER protein coding sequence can be used to generate a variegated population of RISKMARKER or
30 INJURYMARKER fragments for screening and subsequent selection of variants of an RISKMARKER or INJURYMARKER protein. In one embodiment, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of a

RISKMARKER or INJURYMARKER coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA that can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by
5 treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal and internal fragments of various sizes of the RISKMARKER or INJURYMARKER protein.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene
10 products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of RISKMARKER or INJURYMARKER proteins. The most widely used techniques, which are amenable to high throughput analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting
15 library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique that enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify RISKMARKER or INJURYMARKER variants (Arkin and Yourvan (1992)
20 PNAS 89:7811-7815; Delgrave *et al.* (1993) Protein Engineering 6:327-331).

ANTI-RISKMARKER AND ANTI-INJURYMARKER ANTIBODIES

An isolated RISKMARKER or INJURYMARKER protein, or a portion or fragment thereof, can be used as an immunogen to generate antibodies that bind RISKMARKER or INJURYMARKER using standard techniques for polyclonal and monoclonal antibody
25 preparation. The full-length RISKMARKER or INJURYMARKER protein can be used or, alternatively, the invention provides antigenic peptide fragments of RISKMARKER or INJURYMARKER for use as immunogens. The antigenic peptide of RISKMARKER or INJURYMARKER comprises at least 8 amino acid residues of the amino acid sequence encoded by a nucleic acid comprising the nucleic acid sequence shown in RISKMARKER 1-8
30 and INJURYMARKER 1-10 and encompasses an epitope of RISKMARKER or INJURYMARKER such that an antibody raised against the peptide forms a specific immune complex with RISKMARKER or INJURYMARKER. Preferably, the antigenic peptide

comprises at least 10 amino acid residues, more preferably at least 15 amino acid residues, even more preferably at least 20 amino acid residues, and most preferably at least 30 amino acid residues. Preferred epitopes encompassed by the antigenic peptide are regions of RISKMARKER or INJURYMARKER that are located on the surface of the protein, *e.g.*,
5 hydrophilic regions. As a means for targeting antibody production, hydropathy plots showing regions of hydrophilicity and hydrophobicity may be generated by any method well known in the art, including, for example, the Kyte Doolittle or the Hopp Woods methods, either with or without Fourier transformation. See, *e.g.*, Hopp and Woods, 1981, Proc. Nat. Acad. Sci. USA 78: 3824-3828; Kyte and Doolittle 1982, J. Mol. Biol. 157: 105-142, each incorporated herein
10 by reference in their entirety.

RISKMARKER or INJURYMARKER polypeptides or derivatives, fragments, analogs or homologs thereof, may be utilized as immunogens in the generation of antibodies that immunospecifically-bind these protein components. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin
15 molecules, *i.e.*, molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, F_{ab} and $F_{(ab)2}$ fragments, and an F_{ab} expression library. Various procedures known within the art may be used for the production of polyclonal or monoclonal antibodies to an RISKMARKER or INJURYMARKER protein sequence, *e.g.*
20 RISKMAKER 1 or RISKMAKER 6-8, or derivatives, fragments, analogs or homologs thereof. Some of these proteins are discussed below.

For the production of polyclonal antibodies, various suitable host animals (*e.g.*, rabbit, goat, mouse or other mammal) may be immunized by injection with the native protein, or a synthetic variant thereof, or a derivative of the foregoing. An appropriate immunogenic
25 preparation can contain, for example, recombinantly expressed RISKMARKER or INJURYMARKER protein or a chemically synthesized RISKMARKER or INJURYMARKER polypeptide. The preparation can further include an adjuvant. Various adjuvants used to increase the immunological response include, but are not limited to, Freund's (complete and incomplete), mineral gels (*e.g.*, aluminum hydroxide), surface active substances
30 (*e.g.*, lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, etc.), human adjuvants such as *Bacille Calmette-Guerin* and *Corynebacterium parvum*, or similar immunostimulatory agents. If desired, the antibody molecules directed against RISKMARKER or INJURYMARKER can be isolated from the mammal (*e.g.*, from the

blood) and further purified by well known techniques, such as protein A chromatography to obtain the IgG fraction.

The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of RISKMARKER or INJURYMARKER. A monoclonal antibody composition thus typically displays a single binding affinity for a particular RISKMARKER or INJURYMARKER protein with which it immunoreacts. For preparation of monoclonal antibodies directed towards a particular RISKMARKER or INJURYMARKER protein, or derivatives, fragments, analogs or homologs thereof, any technique that provides for the production of antibody molecules by continuous cell line culture may be utilized. Such techniques include, but are not limited to, the hybridoma technique (see Kohler & Milstein, 1975 *Nature* 256: 495-497); the trioma technique; the human B-cell hybridoma technique (see Kozbor, *et al.*, 1983 *Immunol Today* 4: 72) and the EBV hybridoma technique to produce human monoclonal antibodies (see Cole, *et al.*, 1985 In: *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). Human monoclonal antibodies may be utilized in the practice of the present invention and may be produced by using human hybridomas (see Cote, *et al.*, 1983. *Proc Natl Acad Sci USA* 80: 2026-2030) or by transforming human B-cells with Epstein Barr Virus *in vitro* (see Cole, *et al.*, 1985 In: *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96).

According to the invention, techniques can be adapted for the production of single-chain antibodies specific to a RISKMARKER or INJURYMARKER protein (see *e.g.*, U.S. Patent No. 4,946,778). In addition, methods can be adapted for the construction of F_{ab} expression libraries (see *e.g.*, Huse, *et al.*, 1989 *Science* 246: 1275-1281) to allow rapid and effective identification of monoclonal F_{ab} fragments with the desired specificity for a RISKMARKER or INJURYMARKER protein or derivatives, fragments, analogs or homologs thereof. Non-human antibodies can be "humanized" by techniques well known in the art. See *e.g.*, U.S. Patent No. 5,225,539. Antibody fragments that contain the idiotype to a RISKMARKER or INJURYMARKER protein may be produced by techniques known in the art including, but not limited to: (i) an $F_{(ab)2}$ fragment produced by pepsin digestion of an antibody molecule; (ii) an F_{ab} fragment generated by reducing the disulfide bridges of an $F_{(ab)2}$ fragment; (iii) an F_{ab} fragment generated by the treatment of the antibody molecule with papain and a reducing agent and (iv) F_v fragments.

Additionally, recombinant anti-RISKMARKER or INJURYMARKER antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and

non-human portions, which can be made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in PCT International Application No. PCT/US86/02269; European Patent Application No. 184,187; European Patent Application No. 171,496; European Patent Application No. 173,494; PCT International Publication No. WO 86/01533; U.S. Pat. No. 4,816,567; European Patent Application No. 125,023; Better *et al.* (1988) *Science* 240:1041-1043; Liu *et al.* (1987) *PNAS* 84:3439-3443; Liu *et al.* (1987) *J Immunol.* 139:3521-3526; Sun *et al.* (1987) *PNAS* 84:214-218; Nishimura *et al.* (1987) *Cancer Res* 47:999-1005; Wood *et al.* (1985) *Nature* 314:446-449; Shaw *et al.* (1988) *J Natl Cancer Inst.* 80:1553-1559; Morrison (1985) *Science* 229:1202-1207; Oi *et al.* (1986) *BioTechniques* 4:214; U.S. Pat. No. 5,225,539; Jones *et al.* (1986) *Nature* 321:552-525; Verhoeyan *et al.* (1988) *Science* 239:1534; and Beidler *et al.* (1988) *J Immunol* 141:4053-4060.

In one embodiment, methods for the screening of antibodies that possess the desired specificity include, but are not limited to, enzyme-linked immunosorbent assay (ELISA) and other immunologically-mediated techniques known within the art. In a specific embodiment, selection of antibodies that are specific to a particular domain of a RISKMARKER or INJURYMARKER protein is facilitated by generation of hybridomas that bind to the fragment of a RISKMARKER or INJURYMARKER protein possessing such a domain. Antibodies that are specific for one or more domains within a RISKMARKER or INJURYMARKER protein, *e.g.*, domains spanning the above-identified conserved regions of RISKMARKER or INJURYMARKER family proteins, or derivatives, fragments, analogs or homologs thereof, are also provided herein.

Anti-RISKMARKER or anti-INJURYMARKER antibodies may be used in methods known within the art relating to the localization and/or quantitation of a RISKMARKER or INJURYMARKER protein (*e.g.*, for use in measuring levels of the RISKMARKER or INJURYMARKER protein within appropriate physiological samples, for use in diagnostic methods, for use in imaging the protein, and the like). In a given embodiment, antibodies for RISKMARKER or INJURYMARKER proteins, or derivatives, fragments, analogs or homologs thereof, that contain the antibody derived binding domain, are utilized as pharmacologically-active compounds [hereinafter "Therapeutics"].

An anti-RISKMARKER or INJURYMARKER antibody (*e.g.*, monoclonal antibody) can be used to isolate RISKMARKER or INJURYMARKER by standard techniques, such as affinity chromatography or immunoprecipitation. An anti-RISKMARKER or

INJURYMARKER antibody can facilitate the purification of natural RISKMARKER or INJURYMARKER from cells and of recombinantly produced RISKMARKER or INJURYMARKER expressed in host cells. Moreover, an anti-RISKMARKER or INJURYMARKER antibody can be used to detect RISKMARKER or INJURYMARKER protein (*e.g.*, in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the RISKMARKER or INJURYMARKER protein.

Anti-RISKMARKER or INJURYMARKER antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, *e.g.*, to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (*i.e.*, physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

RISKMARKER RECOMBINANT VECTORS AND HOST CELLS

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding RISKMARKER protein, *e.g.*, RISKMARKER 1, or RISKMARKER 6-8, or derivatives, fragments, analogs or homologs thereof. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a linear or circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein

as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, that is operatively linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner that allows for expression of the nucleotide sequence (e.g., in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel; Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, Calif. (1990). Regulatory sequences include those that direct constitutive expression of a nucleotide sequence in many types of host cell and those that direct expression of the nucleotide sequence only in certain host cells (e.g., tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (e.g., RISKMARKER proteins, mutant forms, fusion proteins, etc.).

The recombinant expression vectors of the invention can be designed for expression of RISKMARKER in prokaryotic or eukaryotic cells. For example, RISKMARKER can be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, Calif. (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve
5 three purposes: (1) to increase expression of recombinant protein; (2) to increase the solubility of the recombinant protein; and (3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification
10 of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith and Johnson (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.) that fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

15 Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amrann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990) 60-89).

One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant
20 protein. See, Gottesman, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990) 119-128. Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (Wada *et al.*, (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention
25 can be carried out by standard DNA synthesis techniques.

In another embodiment, the RISKMARKER expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, (1987) *EMBO J* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.*, (1987) *Gene* 54:113-123), pYES2 (Invitrogen Corporation, San Diego, Calif.),
30 and picZ (Invitrogen Corp, San Diego, Calif.).

Alternatively, RISKMARKER can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect

cells (*e.g.*, SF9 cells) include the pAc series (Smith *et al.* (1983) *Mol Cell Biol* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors
5 include pCDM8 (Seed (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.* (1987) *EMBO J* 6: 187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells. See, *e.g.*, Chapters 16
10 and 17 of Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (*e.g.*, tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific
15 regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.* (1987) *Genes Dev* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv Immunol* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J* 8:729-733) and immunoglobulins (Banerji *et al.* (1983) *Cell* 33:729-740; Queen and
20 Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle (1989) *PNAS* 86:5473-5477), pancreas-specific promoters (Edlund *et al.* (1985) *Science* 230:912-916), and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S. Pat. No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, *e.g.*, the murine hox
25 promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the β -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev* 3:537-546).

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner that allows
30 for expression (by transcription of the DNA molecule) of an RNA molecule that is antisense to RISKMARKER mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen that direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen that direct constitutive, tissue specific or cell type specific

expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub *et al.*, "Antisense RNA as a
5 molecular tool for genetic analysis," Reviews--Trends in Genetics, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer
10 not only to the particular subject cell but also to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, RISKMARKER
15 protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and
20 "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (Molecular Cloning: A Laboratory Manual. 2nd ed., Cold Spring Harbor Laboratory, Cold
25 Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that
30 encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Various selectable markers include those that confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be introduced into a host cell on the same vector as that encoding

RISKMARKER or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (*i.e.*, express) an RISKMARKER protein, *e.g.* RISKMARKER 1, or RISKMARKER 6-8. Accordingly, the invention further provides methods for producing RISKMARKER protein using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding RISKMARKER has been introduced) in a suitable medium such that RISKMARKER protein is produced. In another embodiment, the method further comprises isolating RISKMARKER from the medium or the host cell.

PHARMACEUTICAL COMPOSITIONS

The RISKMARKER nucleic acid molecules, RISKMARKER proteins, and anti-RISKMARKER or anti-INJURYMARKER antibodies (also referred to herein as "active compounds") of the invention, and derivatives, fragments, analogs and homologs thereof, can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Suitable carriers are described in the most recent edition of Remington's Pharmaceutical Sciences, a standard reference text in the field, which is incorporated herein by reference. Preferred examples of such carriers or diluents include, but are not limited to, water, saline, finger's solutions, dextrose solution, and 5% human serum albumin. Liposomes and non-aqueous vehicles such as fixed oils may also be used. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*,

intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (*e.g.*, a RISKMARKER protein or anti-RISKMARKER or INJURYMARKER antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are

prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art. The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable,

biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal
5 suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage
10 unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique
15 characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example,
20 intravenous injection, local administration (see U.S. Pat. No. 5,328,470) or by stereotactic injection (see *e.g.*, Chen *et al.* (1994) *PNAS* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, *e.g.*,
25 retroviral vectors, the pharmaceutical preparation can include one or more cells that produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

30 **KITS AND NUCELIC ACID COLLECTIONS FOR IDENTIFYING RISKMARKER AND INJURYMARKER NUCLEIC ACIDS**

In another aspect, the invention provides a kit useful for examining hepatotoxicity of agents. The kit can include nucleic acids that detect two or more RISKMARKER or

INJURYMARKER sequences. In preferred embodiments, the kit includes reagents which detect 3, 4, 5, 6, 8, 10, 12, 15, or all of the RISKMARKER or INJURYMARKER nucleic acid sequences.

5 The invention also includes an isolated plurality of sequences which can identify one or more RISKMARKER or INJURYMARKER responsive nucleic acid sequences. The kit or plurality may include, *e.g.*, sequence homologous to RISKMARKER or INJURYMARKER nucleic acid sequences, or sequences which can specifically identify one or more RISKMARKER or INJURYMARKER nucleic acid sequences.

OTHER EMBODIMENTS

10 It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

WE CLAIM:

1. A method of screening a test agent for hepatotoxicity, the method comprising;
 - (a) providing a test cell population comprising a cell capable of expressing one or more nucleic acid sequences selected from the group consisting of RISKMARKER 1-8 and INJURYMARKER 1-10;
 - (b) contacting the test cell population with a test agent;
 - (c) measuring expression of one or more of the nucleic acid sequences in the test cell population;
 - (d) comparing the expression of the nucleic acid sequences in the test cell population to the expression of the nucleic acid sequences in a reference cell population comprising at least one cell whose exposure status to a hepatotoxic agent is known; and
 - (e) identifying a difference in expression levels of the RISKMARKER or INJURYMARKER sequences, if present, in the test cell population and reference cell population,thereby screening said test agent for hepatotoxicity.
2. The method of claim 1, wherein said hepatotoxicity comprises idiosyncratic hepatotoxicity.
3. The method of claim 2, wherein the method comprises comparing the expression of one or more nucleic acid sequences selected from the group consisting of RISKMARKER 1-8.
4. The method of claim 2, wherein the method comprises comparing the expression of one or more nucleic acid sequences selected from the group consisting of INJURYMARKER 1-10.
5. The method of claim 1, wherein the method comprises comparing the expression of 6 or more of the nucleic acid sequences.
6. The method of claim 1, wherein the expression of the nucleic acid sequences in the test cell population is decreased as compared to the reference cell population.

7. The method of claim 1, wherein the expression of the nucleic acid sequences in the test cell population is increased as compared to the reference cell population.
8. The method of claim 1, wherein the test cell population is provided in vitro.
9. The method of claim 1, wherein the test cell population is provided ex vivo from a mammalian subject.
10. The method of claim 1, wherein the test cell population is provided in vivo in a mammalian subject.
11. The method of claim 1, wherein the test cell population is derived from a human or rodent subject.
12. The method of claim 1, wherein the test cell population includes a hepatocyte.
13. The method of claim 1, wherein said test agent is an idiosyncratic hepatotoxic agent.
14. The method of claim 1, wherein said test agent is a non-steroidal anti-inflammatory drug (NSAID).
15. The method of claim 3, wherein said hepatotoxic agent is a NSAID.
16. The method of claim 15, wherein said NSAID is a NSAID classified as having a low risk of hepatotoxicity, and wherein said test agent is identified as having a low risk of hepatotoxicity if no qualitative difference in expression levels is identified in step (e).
17. The method of claim 16, wherein said difference in expression levels is determined by comparing expression transformation eigenvectors for said test cell and reference cell populations.
18. The method of claim 16, wherein said NSAID is selected from the group consisting of Benoxaprofen, Bromfenac, Diclofenac, Phenylbutazone, and Sulindac.

19. The method of claim 18, wherein said NSAID is selected from the group consisting of Benoxaprofen, Phenylbutazone, and Sulindac.
20. The method of claim 15, wherein said NSAID is a NSAID classified as having a very low risk of hepatotoxicity, and wherein said test agent is identified as having a very low risk of hepatotoxicity if no qualitative difference in expression levels is identified in step (e).
21. The method of claim 20, wherein said difference in expression levels is determined by comparing expression transformation eigenvectors for said test cell and reference cell populations.
22. The method of claim 20, wherein said NSAID is selected from the group consisting of Etodolac, Fenoprofen, Flurbiprofen, Ibuprofen, Indomethacin, Ketoprofen, Meclofenamate, Mefenamic Acid, Nabumetone, Naproxen, Oxaprozin, Piroxicam, Suprofen, Tenoxicam, Tolmentin, and Zomepirac.
23. The method of claim 22, wherein said NSAID is selected from the group consisting of Flurbiprofen, Oxaprozin, and Tenoxicam.
24. The method of claim 15, wherein said NSAID is a NSAID classified as having an overdose risk of hepatotoxicity, and wherein said test agent is identified as having an overdose risk of hepatotoxicity if no qualitative difference in expression levels is identified in step (e).
25. The method of claim 24, wherein said difference in expression levels is determined by comparing expression transformation eigenvectors for said test cell and reference cell populations.
26. The method of claim 25, wherein said NSAID is selected from the group consisting of Acetaminophen, Acetylsalicylic acid, and Phenacetin.
27. The method of claim 4, wherein said hepatotoxic agent is a NSAID.

28. The method of claim 27, wherein said NSAID is a NSAID classified as inducing hepatocellular damage, and wherein said test agent is identified as likely to induce hepatocellular damage if no qualitative difference in expression levels is identified in step (e).
29. The method of claim 28, wherein said difference in expression levels is determined by comparing expression transformation eigenvectors for said test cell and reference cell populations.
30. The method of claim 27, wherein said NSAID is selected from the group consisting of Acetaminophen, Flurbiprofen, and Ketoprofen.
31. The method of claim 27, wherein said NSAID is a NSAID classified as inducing cholestasis, and wherein said test agent is identified as likely to induce cholestasis if no qualitative difference in expression levels is identified in step (e).
32. The method of claim 31, wherein said difference in expression levels is determined by comparing expression transformation eigenvectors for said test cell and reference cell populations.
33. The method of claim 30, wherein said NSAID is selected from the group consisting of Benoxaprofen, Nabumetone, and Sulindac.
34. The method of claim 27, wherein said NSAID is a NSAID classified as inducing elevated transaminase level, and wherein said test agent is identified as likely to induce elevated transaminase level if no qualitative difference in expression levels is identified in step (e).
35. The method of claim 34, wherein said difference in expression levels is determined by comparing expression transformation eigenvectors for said test cell and reference cell populations.

36. The method of claim 34, wherein said NSAID is selected from the group consisting of Zomepirac, Mefenamic acid, and Tenoxicam.
37. A method of assessing the hepatotoxicity of a test agent in a subject, the method comprising:
- (a) providing from the subject a test cell population comprising a cell capable of expressing one or more nucleic acid sequences selected from the group consisting of RISKMARKER 1-8 and INJURYMARKER 1-10;
 - (b) contacting the test cell population with a test agent;
 - (c) measuring expression of one or more of the nucleic acid sequences in the test cell population; and
 - (d) comparing the expression of the nucleic acid sequences in the test cell population to the expression of the nucleic acid sequences in a reference cell population comprising at least one cell whose exposure status to a hepatotoxic agent is known;
 - (e) identifying a difference in expression levels of the nucleic acid sequences, if present, in the test cell population and the reference cell population, thereby assessing the hepatotoxicity of the test agent in the subject.
38. The method of claim 37, wherein said hepatotoxicity comprises idiosyncratic hepatotoxicity.
39. The method of claim 38, wherein the method comprises comparing the expression of one or more nucleic acid sequences selected from the group consisting of RISKMARKER 1-8.
40. The method of claim 38, wherein the method comprises comparing the expression of one or more nucleic acid sequences selected from the group consisting of INJURYMARKER 1-10.
41. The method of claim 37, wherein the expression of the nucleic acid sequences in the test cell population is increased as compared to the reference cell population.

42. The method of claim 37, wherein the expression of the nucleic acid sequences in the test cell population is increased as compared to the reference cell population.
43. The method of claim 37, wherein said subject is a human or rodent.
44. The method of claim 37, wherein the test cell population is provided ex vivo from said subject.
45. The method of claim 37, wherein the test cell population is provided in vivo from said subject.
46. The method of claim 37, wherein said test agent is a non-steroidal anti-inflammatory drug (NSAID).
47. The method of claim 37, wherein said hepatotoxic agent is a NSAID.
48. An isolated nucleic acid comprising a nucleic acid sequence selected from the group consisting of a RISKMARKER 1 nucleic acid, a RISKMARKER 6-8 nucleic acid, and their complements.
49. A vector comprising the nucleic acid of claim 48.
50. A cell comprising the vector of claim 49.
51. A pharmaceutical composition comprising the nucleic acid of claim 48.
52. A polypeptide encoded by the nucleic acid of claim 48.
53. An antibody which specifically binds to the polypeptide of claim 52.

54. A kit which detects two or more of the nucleic acid sequences selected from the group consisting of RISKMARKER 1, and RISKMARKER 6-8.
55. An array which detects one or more of the nucleic acid selected from the group consisting of RISKMARKER 1, and RISKMARKER 6-8.
56. A plurality of nucleic acid comprising one or more of the nucleic acid selected from the group consisting of RISKMARKER 1, and RISKMARKER 6-8.